

# Preparation of Wound Healing Gel and its Evaluation

**Mr. Pranav T. Pandhare<sup>1</sup> and Mr. Pramod B. Chikkodi<sup>2</sup>**

UG Student, Department of Pharmaceutics<sup>1</sup>

HOD, Department of Pharmaceutics<sup>2</sup>

Nootan College of Pharmacy, Kavathe Mahankal, Sangli, Maharashtra, India

**Abstract:** In this present study, we have prepared herbal gel for wound management and its batches. currently issues related to the wound and its repairing mechanism time taking process. Required material for the preparation of gel we taken by the common process we have prepared gel. By different method the formulation was prepared and further its evaluation test was performed. Out of this batch last HG were found effective and best as comparative other batches. Henceforth as per our need herbal gel formulation were prepared and its ideal evaluation process were carried out.

**Keywords:** Herbal gel, Wound healing, Evaluation

## REFERENCES

- [1]. Babu, M., Gnanamani, A., Radhakrishnan, N., and Priya, K. (2002) Healing potential of Datura alba on burn wounds in albino rats. J. Ethnopharmacology, 83, pp. 193-199.
- [2]. Perumal, S.R., Ignacimuthu, S., and Sen, A. (1998) Screening of 34 Indian medicinal plants for antibacterial properties. Ethnopharmacol., 62(2), pp. 173-182.
- [3]. Srikanth, D., Shenoy, R., and Rao, C.M. (2008) The effects of topical (gel) astemizole and terfenadine on wound healing. Indian J. Pharmacology, 40(4), pp. 170174.
- [4]. Liu, M., Dai, Y., Li, Y., and Luo, Y. (2008) Madecassoside isolated from centalla asiatica herbs facilitates burn wound healing in mice. Planta Med., 74(8), pp. 809-815.
- [5]. Vure P, Dorl AK, Evaluation of ghee-based formulation for wound healing activity, J Ethnopharmacol, 2006, 107, 38-47.
- [6]. Bhat S, Shankrappa J, Kumar G, Formulation and evaluation of polyherbal wound treatments, J Pharm Sci, 2007, 2, 11-17.
- [7]. Ramane S, Syed V, Biyani K, Evaluation of wound healing activity of polyherbal gel a novel herbal formulation, Int J Res Pharm Biomed Sci, 2011, 3, 39-42
- [8]. Kikwai L, Babu RJ, Prado RA, Kolot A, Armstrong CA, Ansel JC et al. In vitro and in vivo evaluation of topical formulations of spantide II. AAPS PharmSciTech 2005; 6 (4): E562-72.
- [9]. Ali Mohammad A., Hamid S. Mohmad K.; The effect of Teucrium Polium Honey on the Wound Healing and tensile strength in rat.; Iranian Journal of Basic Medical Science, 2011; 14(6): 499-505.
- [10]. Wadankar G.D., Malode S.N.; Traditionally Used Medicinal Plants for Wound Healing in the Washim District, Maharashtra (India).; International Journal of Pharm Tech Research, 2011; 3(4): 2080-2084
- [11]. Dr. K.R. Khandelwal; Practical Pharmacognosy Technniqes And Experiments; Nirali Prakashan, 2011; 23: 13-23.14.
- [12]. Karade Preeti G., Shah Rohit R.; Formulation and evaluation of celecoxib gel; Journal of Drug Delivery and therapeutics, 2012; 3: 132-135