



IJARSCT

International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Impact Factor: 7.53

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 4, Issue 3, March 2024

ISSN (Online) 2581-9429

The One Pot Three Component Synthesis of 2-Amino-4H-Chromenes by using Sulfonic Acid-Functionalized Phthalimide

Mahesh Walle

Department of Chemistry

Sundarrao More Arts, Commerce & Science College, Poladpur, Raigad, Maharashtra, India

mahesh.walle@gmail.com

Abstract: Sulfonic acid-functionalized phthalimide efficiently catalyzes the one-pot, three component reaction of an aromatic aldehyde, malonitrile and α or β -naphthol to yield 2-amino-4H-chromenes in very good yields. The catalyst sulfonic acid-functionalized phthalimide (SFP) is reusable and efficient in multicomponent reactions. The reaction is carried out in green approach which is environment friendly with excellent yield.

Keywords: Sulfonic acid-functionalized phthalimide; 2-Amino-4H-chromenes; One-pot reaction

REFERENCES

- [1]. Katritzky, A. R., Rees, C. W., & Drayton, C. J. (1984). Comprehensive heterocyclic chemistry: the structure, reactions, synthesis, and uses of heterocyclic compounds. (*No Title*).
- [2]. Baghernejad, B., Heravi, M. M., & Oskooie, H. A. (2009). A novel and efficient catalyst to one-pot synthesis of 2-amino-4H-chromenes by p-toluenesulfonic acid. *Journal of the Korean Chemical Society*, 53(6), 631-634.
- [3]. Konkoy, C. S., Fick, D. B., Cai, S. X., Lan, N. C., & Keana, J. F. W. (2001). PCT Int. Appl. WO 0075123, 2000. In *Chem. Abstr* (Vol. 134, p. 29313a).
- [4]. Ballini, R., Bosica, G., Conforti, M. L., Maggi, R., Mazzacani, A., Righi, P., & Sartori, G. (2001). Three-component process for the synthesis of 2-amino-2-chromenes in aqueous media. *Tetrahedron*, 57(7), 1395-1398.
- [5]. Heravi, M. M., Baghernejad, B., & Oskooie, H. A. (2008). A Novel and Efficient Catalyst to One-Pot Synthesis of 2-Amino-4H-Chromenes by Methanesulfonic Acid. *Journal of the Chinese Chemical Society*, 55(3), 659-662.
- [6]. Sharghi, H., & Sarvari, M. H. (2003). Synthesis of macrocyclic polyether-diester compounds using Al₂O₃/MeSO₃H (AMA) as a new reagent. *Synthesis*, 2003(06), 0879-0882.
- [7]. Sarvari, M. H. (2005). ZnO/CH₃COCl: A new and highly efficient catalyst for dehydration of aldoximes into nitriles under solvent-free condition. *Synthesis*, 787-790.
- [8]. Sharghi, H., & Sarvari, M. H. (2003). Al₂O₃/MeSO₃H (AMA) as a new reagent with high selective ability for monoesterification of diols. *Tetrahedron*, 59(20), 3627-3633.
- [9]. Niknam, K., & Ali Zolfigol, M. (2006). Alumina-Methanesulfonic Acid (AMA)/NaNO₂ as an Efficient Procedure for the Chemoselectivite N-Nitrosation of Secondary Amines. *Synthetic communications*, 36(16), 2311-2319.
- [10]. Niknam, K., Zolfigol, M. A., Razavian, S. M., & Mohammahpoor-Baltork, I. (2006). Aromatization of 1, 4-Dihydropyridines in the Presence of Methanesulfonic Acid/NaNO₂/Wet SiO₂ under Both Heterogeneous and Solvent Free Conditions. *Journal of heterocyclic chemistry*, 43(1), 199-202.
- [11]. Tian, Y., Meng, X., Duan, J. Y., & Shi, L. (2012). A novel application of methanesulfonic acid as catalyst for the alkylation of olefins with aromatics. *Industrial & engineering chemistry research*, 51(42), 13627-13631.



IJARSCT

International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

IJARSCT

ISSN (Online) 2581-9429

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Impact Factor: 7.53

Volume 4, Issue 3, March 2024

- [12]. Elagamey, A. G. A., & El-Taweel, F. M. A. A. (1990). Nitriles in Heterocyclic Synthesis: Synthesis of Condensed Pyrans. *ChemInform*, 21(49), no-no.
- [13]. Ballini, R., Bosica, G., Conforti, M. L., Maggi, R., Mazzacani, A., Righi, P., & Sartori, G. (2001). Three-component process for the synthesis of 2-amino-2-chromenes in aqueous media. *Tetrahedron*, 57(7), 1395-1398.
- [14]. Babu, T. H., & Perumal, P. T. (2011). A simple Method for the Synthesis of Functionalised Chromenes via Vinylogous Michael Addition of α, α -Dicyanoalkenes on iminocoumarin Derivatives. *Synlett*, 2011(03), 341-344.
- [15]. Maggi, R., Ballini, R., Sartori, G., & Sartorio, R. (2004). Basic alumina catalysed synthesis of substituted 2-amino-2-chromenes via three-component reaction. *Tetrahedron letters*, 45(11), 2297-2299.
- [16]. Jin, T. S., Xiao, J. C., Wang, S. J., & Li, T. S. (2004). Ultrasound-assisted synthesis of 2-amino-2-chromenes with cetyltrimethylammonium bromide in aqueous media. *Ultrasonics Sonochemistry*, 11(6), 393-397.
- [17]. Walle, M., Pansare, D., Khan, T., Pawar, R., Shelke, R., & Ingle, R. (2021). One pot three component synthesis of 2-amino-5-oxo-4, 5-dihydropyrano [3, 2-c] chromene-3-carbonitrile derivatives catalyzed by cobalt doped iron (III) tartarate complex. *Lett App NanoBioSci*, 11(1), 3208-3217.
- [18]. Walle, M., Ingle, R., & Pawar, R. (2021). Efficient and One-pot Synthesis of Tetrahydro [b] Pyran Derivatives Catalyzed by Copper Doped Iron Tartarate. *Journal of Scientific Research*, 65(6).
- [19]. Walle, M. R., Pansare, D. N., Kamble, S. S., Pawar, R. P., & Ingale, R. D. (2019). Synthesis of 1, 8-dioxooctahydroxanthene and 3, 3-arylidene bis (4-hydroxycoumarin) derivatives. *European Chemical Bulletin*, 8(3), 101-104.
- [20]. Ziarani, G. M., Lashgari, N., & Badiei, A. (2015). Sulfonic acid-functionalized mesoporous silica (SBA-Pr-SO₃H) as solid acid catalyst in organic reactions. *Journal of Molecular Catalysis A: Chemical*, 397, 166-191.
- [21]. Heravi, M. M., Hashemi, E., & Azimian, F. (2015). N-Sulfonic acid modified poly (styrene-co-maleic anhydride): an efficient and recyclable solid acid catalyst for the synthesis of a wide range of spiropyrans. *Journal of the Iranian Chemical Society*, 12, 647-653.
- [22]. Moosavi-Zare, A. R., Zolfogol, M. A., Khakyzadeh, V., Böttcher, C., Beyzavi, M. H., Zare, A., ... & Luque, R. (2014). Facile preparation of a nanostructured functionalized catalytically active organosalt. *Journal of Materials Chemistry A*, 2(3), 770-777.
- [23]. Khalafi-Nezhad, A., Panahi, F., Mohammadi, S., & Foroughi, H. O. (2013). A green and efficient procedure for one-pot synthesis of xanthenes and acridines using silica boron-sulfuric acid nanoparticles (SBSANs) as a solid Lewis-protic acid. *Journal of the Iranian Chemical Society*, 10, 189-200.
- [24]. RezaáMoosavi-Zare, A., & AliáZolfogol, M. (2013). Synthesis of 6-amino-4-(4-methoxyphenyl)-5-cyano-3-methyl-1-phenyl-1, 4-dihydropyrano [2, 3-c] pyrazoles using disulfonic acid imidazoliumchloroaluminate as a dual and heterogeneous catalyst. *New Journal of Chemistry*, 37(12), 4089-4094