

# **Valuation of Genotoxic Effect of Aqueous Organic Compound with Allium Sativum [Garlic]**

**Smita S. Kharkale-Bhuyar**

Assistant Professor

Shri Lemdeo Patil Mahavidyalaya, Mandhal, Kuhi, Nagpur, Maharashtra, India

**Abstract:** The current study was deliberate to estimate the genotoxic effect of 1,2,4,5-tetrazin using Allium sativum [Garlic] root chromosomal abnormality analyse. Root tips of Allium sativum were treated with different concentrations of 0.001M, 0.002M, 0.003M, 1,2,4,5-tetrazin with 10% Dimethyl Sulfoxide-Water were used in the current study. After treatment Various types of physiological hydrolyzed, squashed, mitotic index (MI), cytological abnormalities and Chromosomal aberrations rate were scored. Then the results exposed a genotoxicity effect as well as significance reduction in the Mitotic index of toxic effect was found to be dose dependent.

**Keywords:** Allium Sativum., Garlic, Mitotic Index, 1,2,4,5-Tetrazin, Chromosomal Aberrations

## **REFERENCES**

- [1]. Nagpal AK, Grover IS (1994) Genotoxic evaluation of systemic pesticides in Allium cepa. L. Mitotic effects. *The Nucleus*. 37: 99-105.
- [2]. Sharma S, Vig AP, (2012) Genotoxicity of Atrazine, Avenoxan, Diuron and Quizalofop-p-ethyl Herbicides using the Allium cepa Root Chromosomal Aberration Assay. *Terrestrial and Aquatic Environmental Toxicology*, 6(2): 90-95.
- [3]. Mishra GM, Sinha SP (1979) Effect of malathion on mitotically dividing onion (Allium cepa) root tip cells. *Indian Journal of Experimental Biology*. 17:716-717.
- [4]. Mohandas T, Grant WF (1972) Cytogenic effects of 2,4-D and amine triazole in relation to nuclear volume and DNA content in some higher plants. *Canadian Journal of Genetics and Cytology*. 14:773-783.
- [5]. Saxena PN, Chauhan LKS, Chandra S, Gupta SK (2004) Genotoxic effects of diuron contaminated soil on the root meristem cells of Allium sativum: a possible mechanism of chromosome damage. *Toxicology Mechanisms and Methods*. 14: 281– 286.
- [6]. Oladele EO, Odeigah PGC, Taiwo, IA (2013) The genotoxic effect of lead and zinc in cowpea groundnut (*Vigna subterranean*). *African Journal of Environmental Science and Technology*. 7:9-13.
- [7]. Werbel, LM, McNamara DJ, Colbry NL, (1979) Synthesis and Antimalarial Effects of N,N-Dialky - 6 - (sub. phenyl) - 1,2,4,5- tetrazin - 3 amines(1,2), *J. Heterocyclic Chem*, (16): 881.
- [8]. Thompson PE, Werbel LM, Antimalaria Agents, Chemistry and Pharmacology, Academic Press, New York, N.T. 1972; (16): 885.
- [9]. Qian Y, Tiffany-Castiglioni E (2003). Lead-induced endoplasmic reticulum (ER) stress responses in the nervous system. *Neurochemical Research*. 28: 153–162.
- [10]. Wiley PF, The Chemistry of Heterocyclic Compounds, The 1,2,4-Triazines, 1,2,4,5 – Tetrazines and Pentazines. A. Weissberger, Ed., Interscience, New York, 1956; 16-881.
- [11]. Qian Y, Tiffany-Castiglioni E (2003). Lead-induced endoplasmic reticulum (ER) stress responses in the nervous system. *Neurochemical Research*. 28: 153–162.
- [12]. Gul T, Kaymak F, Fulya D, Muranli G (2006). Genotoxic Effects of Avenoxan on Allium cepa L. and Allium sativum L. *Caryologia*. 59: 241-247.
- [13]. Lanzotti V. The analysis of onion and garlic. *J Chromatogr A* 2006;1112:3-22
- [14]. Christofoletti CA, Pedro-Escher J, Fontanetti CS. Assessment of the genotoxicity of two agricultural

- residues after processing by diplopods using the Allium cepa assay. Water Air Soil Pollut 2013;224:1e14.
- [15]. Bhat SA, Singh J, Vig AP. Vermistabilization of sugar beet (*Beta vulgaris L*) waste produced from sugar factory using earthworm *Eisenia fetida*: genotoxic assessment by Allium cepa test Environ Sci Pollut Res Int 2015;22:11236e54.
- [16]. Saxena PN, Gupta SK, Murthy RC. Carbofuran induced cytogenetic effects in root meristem cells of Allium cepa and Allium sativum: a spectroscopic approach for chromosome damage. Pestic Biochem Physiol 2010;96:93e100.
- [17]. Kharkale SS, Bhuyar SS, Agrawal PS and Paliwal L J (2017). The Evaluation of various organic solvent using Allium Cepa Test. European Journal of Biomedical and Pharmaceutical Sciences. 4: 820-823.
- [18]. Xiao D, Singh SV. Diallyl trisulfide, a constituent of processed garlic, inactivates Akt to trigger mitochondrial translocation of BAD and caspase-mediated apoptosis in human prostate cancer cells. Carcinogenesis 2006;27:533-40.
- [19]. Saxena PN, Gupta SK, Murthy RC (2010). Carbofuran inducedcy to genetic effects in root meristem cellsof Allium cepa and Allium sativum: A spectroscopic approach for chromosome damage. Pesticide Biochemistry and Physiology. 96:93–100.
- [20]. Kim JY, Kwon O. Garlic intake and cancer risk: An analysis using the Food and Drug Administration's evidence-based review system for the scientific evaluation of health claims. Am J Clin Nutr 2009;89:257-64.
- [21]. Agarwal SS, Singh VK. Immunomodulators: A review of studies on Indian medicinal plants and synthetic peptides part I; medicinal plants. Proceedings of the Indian National science academy 1999;65:179-204.
- [22]. Pandir D. (2018) Assesment of the Genotoxic Effect of the Diazinon on Root Cells of Allium cepa (L.) Biological and Applied Sciences 61: 1-13
- [23]. Klasterska I, Natrajan AT, Ramel C (1976). An interception of the origin of sub-chromatid aberrations and chromosome stickiness as a category of chromatid aberrations. Hereditas. 83:153-162.
- [24]. Levan A (1938). The effect of colchicine on root mitosis in Allium. Hereditas. 24: 471-486.
- [25]. Gul T, Kaymak F, Fulya D, Muranli G (2006). Genotoxic Effects of Avenoxan on Allium cepa L. and Allium sativum L. Caryologia. 59: 241-247.
- [26]. Liu D, Jiang W, Meng Q, Zou J, Gu J, Zeng M (2009). Cytogenetical and ultrastructural effects of copper on root meristem cells of Allium sativum L. Biocell. 33: 25-32.
- [27]. Amer SM, Farah OR (1974). Cytological effects of pesticides. VI. Effect of insecticide rogor on the mitosis of *Vicia faba* and *Gossypium barbadense*. Cytologia. 39: 507-514.
- [28]. Singh BD, Singh Y, Singh RB, Singh RM, Sarma NDRK, Singh J (1977). Cytogenetic aberrations and morphological changes induced by insecticide treatments of barley seeds. Indian Journal of Experimental Biology. 15: 688-691.
- [29]. Gul T, Kaymak F, Fulya D, Muranli G (2006). Genotoxic Effects of Avenoxan on Allium cepa L. and Allium sativum L. Caryologia. 59: 241-247.
- [30]. Saxena PN, Gupta SK, Murthy RC (2010). Carbofuran inducedcy to genetic effects in root meristem cells of Allium cepa and Allium sativum: Aspectroscopic approach for chromosome damage. Pesticide Biochemistry and Physiology. 96:93–100
- [31]. Levan A (1938). The effect of colchicine on root mitosis in Allium. Hereditas. 24: 471-486. Singh, Scientific Reports 2020, 10:886