

Status of the Freshwater Molluscs from Krishna River, Satara, (MS) India

Santosh P. Supanekar

Department of Zoology

Mahatma Phule Arts Science Commerce College, Panvel, Maharashtra, India

Corresponding author: sankam23@gmail.com

Abstract: A total 18 freshwater mollusc species with all forms and varieties belonging to 04orders, under 10 families. the maximum diversity was found in order Mesogastropoda and minimum was from Basommatophora. The dominant species among these total molluscs belonged to the Unionidae family. The abundance of freshwater molluscs species in the Krishna River is described in an assessed checklist. Molluscs are widely regarded as the most diverse and dominant benthic fauna in both the lentic and lotic zones. Regional populations of malacofauna, on the other hand, may be attributable to the quantity of planktonic mass at different river sites, which reflects molluscs species diversity.

Keywords: Bivalves, Freshwater molluscs, Gastropods, Krishna River.

REFERENCES

- [1]. Bouchet P., Rocroi J. P., Fryda J., Hausdorf B., Ponder W., Valdés A. and Waren A. (2005): Classification and Nomenclature of gastropod families. *Malacologia: International Journal of Malacology*. Hackenheim, Germany: ConchBooks. 47 (1-2):1-397.
- [2]. Wilbur K. M. and Yonge C. M., (1964): *Physiology of Mollusca*, Academic Press, New York, pp: 473.
- [3]. Oehlmann, J.; Schulte-Oehlmann, U. Molluscs as bioindicators. In *Trace Metals and Other Contaminants in the Environment*; Markert, B.A., Breure, A.M., Zechmeister, H.G., Eds.; Elsevier: Amsterdam, The Netherlands, 2003; pp. 577–635.
- [4]. Heino, J.; Muotka, T.; Paavola, R.; Hämäläinen, H.; Koskenniemi, E. Correspondence between regional delineations and spatial patterns in macroinvertebrate assemblages of boreal headwater streams. *J. N. Am. Benthol. Soc.* 2002, 21, 397–413.
- [5]. Poff, N.L. Landscape filters and species traits: Towards mechanistic understanding and prediction in stream ecology. *J. N. Am. Benthol. Soc.* 1997, 16, 19
- [6]. Ruppert, E.E., Fox, R.S., Barnes, R.D., 2004. *Invertebrate Zoology*. Thomson Learning, Inc, Belmont, California.
- [7]. Subba Rao, N.V. and A. Dey (1989). Freshwater Molluscs in Aquaculture, pp. 225-232. In: *Handbook of Freshwater Molluscs of India*. Zoological Survey of India, Calcutta, 289pp.
- [8]. Vaughn CC, Gido KB, Spooner DE (2004) Ecosystem processes performed by unionid mussels in stream mesocosms: species roles and effects of abundance. *Hydrobiologia*, 527: 35-47
- [9]. János Salánki I, Anna Farkas, Tamara Kamardina, Katalin S Rózsa (2003) Molluscs in biological monitoring of water quality, *Toxicol Lett* 11;140-141:403-10. doi: 10.1016/s0378-4274(03)00036-5.
- [10]. Strong, E.E., Gargominy, O., Ponder, W.F. and Bouchet, P. (2008). Global diversity of gastropods (Gastropoda; Mollusca) in freshwater., *Hydrobiologia* 595:149–166.
- [11]. Subba Rao N (1993) Freshwater molluscs of India. In: Rao KS (Ed.) *Recent advances in freshwater biology* 2. Anmol Publications, New Delhi, India, 187–202
- [12]. Ramakrishna, Dey A (2007) *Handbook on Indian freshwater molluscs*. Zoological Survey of India, Kolkata, India, 399 pp.

- [13]. Sankarappan A, Chellapandian B, Vimalanathan A, Mani K, Sundaram D, Muthukalingan K (2015) Vector ecology of human schistosomiasis in south India and description of a new species of the genus Ferrissia (Mollusca: Gastropoda: Planorbidae). *Journal of Vector Borne Diseases* 52 (3): 201–207
- [14]. Molluscabase (2021) <https://www.molluscabase.org>. Accessed on: 2021-02-15.
- [15]. Patil S. G. (2003) Occurance of freshwater Bivalve (Bivalvia:Unionidae) in Pusad. Yavatmal. District. Maharashtra.Zoos Print. Journal;18(a):11-95.
- [16]. Johnpaul A., Ragunathan M. B. and Selvanayagam M., (2010): Population dynamics of freshwater molluscs in lentic ecosystems in and around Chennai., Rec. Res. Sci. Tech. 2: 80-86.
- [17]. Rao S. and Dey A., (1986): Freshwater molluscs of Misoram, J. Hydrobiol. 2: 25-32.
- [18]. Patil S.G. and Talmale S.S. (2005). A Checklist of Land and Freshwater Mollusca of Maharashtra State. Zoo's PrintJ. 20(6): 1912-1913.
- [19]. Dutta S.P. and Malhotra Y.R. (1986). Seasonal variation in macro benthic fauna of Gandigarh stream (Miran Sahib),Jamm. Indian J. Eco. 13: 138-145.
- [20]. Verma A.K. and Saksena D.N. (2010). Impact of pollution on sewage collecting river Kalpi (Morar) Gwalior (M.P.)with special reference to water quality and Macrozoobenthic Fauna. Asian J. Exp. Bio. Sci. (1):155-161.