

Plant Latex- A Concise Review on the Exudate

Sanjukta Rajhans¹ and Himanshu Pandya²

Department of Botany, Bioinformatics, Climate Change Impacts Management
Gujarat University, School of Sciences, Ahmedabad, Gujarat, India
sanjuktarajhans@gmail.com

Abstract: In this review article an effort has been made to provide basic knowledge about plant latex. In this paper the origin and evolutionary aspects of latex has been mentioned. The geographical distribution and classification along with the physical and anatomical traits have also been brought into limelight. Moreover, the presence of phytochemicals, proteins and variation of the content of rubber in latex has been referred in the paper respectively. This paper would be of great help in gaining deeper knowledge about the plant latex for future researches.

Keywords: Laticifers, Rubber, Diversity, Secretion

REFERENCES

- [1]. Bushman, B. S., Scholte, A. A., Cornish, K., Scott, D. J., Brichta, J. L., Vederas, J. C., Ochoa, O., Michelmore, R. W., Shintani, D. K., & Knapp, S. J. (2006). Identification and comparison of natural rubber from two *Lactuca* species. *Phytochemistry*, 67(23), 2590-2596.
- [2]. Cai, X., Li, W., & Yin, L. (2009). Ultrastructure and cytochemical localization of acid phosphatase of laticifers in *Euphorbia kansui* Liou. *Protoplasma*, 238(1-4), 3-10.
- [3]. Carter, C. A., Forney, R. W., Gray, E. A., Gehring, A. M., Schneider, T. L., Young, D. B., Lovett Jr., C. M., Scott, L., Messer, A. C. & Richardson, D. P. (1997). Toxicarioside A. A new cardenolide isolated from *Antiaristoxicaria* latex-derived dart poison. Assignment of the ¹H- and ¹³C-NMR shifts for an antiarigenin aglycone. *Tetrahedron*, 53(40), 13557-13566.
- [4]. Chavan, B. A., Hundiwale, J. C., & Patil, A. V. (2015). Plant Latex: An Inherent Spring of Pharmaceuticals. *World Journal of Pharmacy and Pharmaceutical Sciences*, 4(4), 1781-1796.
- [5]. Dussourd, D. E. & Denno, R. F. (1991). Deactivation of Plant Defense: Correspondence Between Insect Behavior and Secretory Canal Architecture. *Ecology*, 72(4), 1383-1396.
- [6]. Dussourd, D. E. & Hoyle, A. M. (2000). Poisoned plusiines: toxicity of milkweed latex and cardenolides to some generalist caterpillars. *Chemoecology*, 10(1), 11-16.
- [7]. Dussourd, D. E. (1993). Foraging with finesse caterpillar adaptations for circumventing plant defense. In N. E. Stamp., & T. M. Casey (Eds.), *Caterpillars: Ecological and Evolutionary Constraints on Foraging* (pp. 92-131). Chapman & Hall.
- [8]. Dussourd, D. E. (1995). Entrapment of Aphids and Whiteflies in Lettuce Latex. *Annals of The Entomological Society of America*, 88(2), 163-172.
- [9]. Farrell, B. D., Dussourd, D. E., & Mitter, C. (1991). Escalation of plant defense: do latex and resin canals spur plant diversification? *The American Naturalist*, 138(4), 881-900.
- [10]. Felpin, F. X., & Lebreton, J. (2004). History, Chemistry and biology of alkaloids from *Lobelia inflata*. *Tetrahedron*, 60(45), 10127-10153.
- [11]. Gershenson, J., & Croteau, R. (1991). Terpenoids. In G. A. Rosenthal & M. R. Berenbaum (Eds.), *Herbivores: Their Interactions with Secondary Plant Metabolites* (2nd ed., pp. 615-219). Academic Press.
- [12]. Gidrol, X., Chrestin, H., Tan, H. L. and Kush, A. (1994). Hevein, a Lectin-like Protein from *Hevea brasiliensis* (Rubber Tree) Is Involved in the Coagulation of Latex. *The Journal of Biological Chemistry*, 269(12), 9278-9283.
- [13]. Hagel, J. M., Yeung, E. C. & Facchini, P. J. (2008). Got milk? The secret life of laticifers. *Trends in Plant Science*, 13(12), 631-639.
- [14]. Hartmann, T. (1991). Alkaloids. In G. A. Rosenthal & M. R. Berenbaum (Eds.), *Herbivores: Their Interactions with Secondary Plant Metabolites* (2nd ed., pp. 79-121). Academic Press.

- [15]. Hirayama, C., Konno, K., Wasano, N., & Nakamura, M. (2007). Differential effects of sugar-mimic alkaloids in mulberry latex on sugar metabolism and disaccharides of Eri and domesticated silkworms: enzymatic adaptation of *Bombyx mori* to mulberry defense. *Insect Biochemistry and Molecular Biology*, 37(12), 1348-1358.
- [16]. Itenov, K., Mølgaard, P., & Nyman, U. (1999). Diurnal fluctuations of the alkaloid concentration in latex of poppy *Papaver somniferum* due to day-night fluctuations of the latex water content. *Phytochemistry*, 52(7), 1229-1234.
- [17]. Jia, S., Li, Y., Dai, X., Li, X., Zhou, Y., Xu, Y., & Wang, H. (2020). Physiological adaptations to sugar-mimic alkaloids: Insights from *Bombyx mori* for long-term adaptation and short-term response. *Ecology and Evolution*, 10(18), 9682-9695.
- [18]. Konno, K. (2011). Plant latex and other exudates as plant defense systems: roles of various defense chemicals and proteins contained therein. *Phytochemistry*, 72(13), 1510-1530.
- [19]. Konno, K., Hirayama, C., Nakamura, M., Tateishi, K., Tamura, Y., Hattori, M., & Kohno, K. (2004). Papain protects papaya trees from herbivorous insects: role of cysteine proteases in latex. *The Plant Journal: For Cell and Molecular Biology*, 37(3), 370-378.
- [20]. Lewinsohn, T. M. (1991). The geographical distribution of plant latex. *Chemoecology*, 2(1), 64-68.
- [21]. Lynn, K. R. & Clevette-Radford, N. A. (1987). Biochemical Properties of Latices from the Euphorbiaceae. *Phytochemistry*, 26(4), 939-944.
- [22]. Metcalfe, C. R. (1967). Distribution of latex in the plant kingdom. *Economic botany*, 21(2), 115-127.
- [23]. Mishra, A. & Parida, S. (2020). Latex of Plants: Wonders of Nature for its therapeutic potentials and a valuable resource towards new drug development. *International Journal of Botany Studies*, 5(6), 334-338.
- [24]. Mooibroek, H. & Cornish, K. (2000). Alternative sources of natural rubber. *Applied Microbiology and Biotechnology*, 53(4), 355-365.
- [25]. Noack, E. A., Crea, A. E. G., & Falsone, G. (1980). Inhibition of mitochondrial oxidative phosphorylation by 4-deoxyphorbol triester, a poisonous constituent of the latex sap of *Euphorbia biglandulosa* Desf. *Toxicicon*, 18(2), 165-174.
- [26]. Oppel, C. B., Dussourd, D. E., & Garimella, U. (2009). Visualizing a Plant Defense and Insect Counterploy: Alkaloid Distribution in *Lobelia* Leaves Trenched by a Plusidine Caterpillar. *Journal of Chemical Ecology*, 35(6), 625-634.
- [27]. Rasmann, S., Johnson, M. D. & Agrawal, A. A. (2009). Induced responses to herbivory and jasmonate in three milkweed species. *Journal of Chemical Ecology*, 35(11), 1326-1334.
- [28]. Rees, S. B., & Harborne, J. B. (1985). The role of sesquiterpene lactones and phenolics in the chemical defense of the chicory plant. *Phytochemistry*, 24(10), 2225-2231.
- [29]. Reitsma, J. M. (1988). Forest vegetation of Gabon. *Tropenbos Technical Series*, 1(1), 5-142.
- [30]. Schmeller, T., Latz-Brüning, B., & Wink, M. (1997). Biochemical activities of berberine, palmatine and sanguinarine mediating chemical defense against microorganisms and herbivores. *Phytochemistry*, 44(2), 257-266.
- [31]. Seiber, J. N., Nelson, C. J., & Lee, S. M. (1982). Cardenolides in the latex and leaves of seven *Asclepias* species and *Calotropis procera*. *Phytochemistry*, 21(9), 2343-2348.
- [32]. Sessa, R. A., Bennett, M. H., Lewis, M. J., Mansfield, J. W., & Beale, M. H. (2000). Metabolite profiling of sesquiterpene lactones from *Lactuca* species. Major latex components are novel oxalate and sulfate conjugates of lactucin and its derivatives. *The Journal of Biological Chemistry*, 275(35), 26877-26884.
- [33]. Snook, M. E., Data, E. S., & Kays, S. J. (1994). Characterization and Quantification of Hexadecyl, Octadecyl and Eicosyl Esters of p-Coumaric Acid in the Vine and Root Latex of Sweetpotato [*Ipomoea batatas* (L.) Lam.]. *Journal of Agricultural and Food Chemistry*, 42(11), 2589-2595.
- [34]. Tomè, F., & Colombo, M. L. (1995). Distribution of alkaloids in *Chelidonium majus* and factor affecting their accumulation. *Phytochemistry*, 40(1), 37-39.
- [35]. Waldhoer, M., Bartlett, S. E., & Whistler, J. L. (2004). Opioid receptors. *Annual Review of Biochemistry*, 73(1), 953-990.
- [36]. Wasano, N., Konno, K., Nakamura, M., Hirayama, C., Hattori, M., & Tateishi, K. (2009). A unique latex protein, MLX56, defends mulberry trees from insects. *Phytochemistry*, 70(7), 880-888.

- [37]. Zhou, Y. F., & Liu, W. Z. (2010). Laticiferous canal formation in fruits of *Decaisneafargesii*: a programmed cell death process? *Protoplasma*, 248(4), 683-694.