

A Review Paper on Various Plants are used in Various Types of Diseases

Shinge Tejaswini Sajjan

Nootan College of Pharmacy, Kavthe Mahankal, Maharashtra, India

Abstract: Medicinal plant are mostly used in healthcare in the whole world. The medicinal plant. are Effective in production of medicine . In this paper we can seen the role and usefulness of medicinal plants and approaches to diseases prevention.

Keywords: Medicinal Plants, Prevention, Natural Source

REFERENCES

- [1]. Ms. Mary Vineela .PMSc (N) Lecturer C H N,Naryana College of Nursing, Nellore, Chemical composition of Tulsi, Benefits of holy basil(tulsi) leafs.
- [2]. Das SK and Vasudevan DM. Tulsi: The Indian holy power plant. Natural Product Radiance. 2006;5:279-83.
- [3]. Triveni, Kuldeep Kumar, Amit Kumar Singh, Rahul Kumar, Vaishnavee Gupta, Kishu Tripathi* Smt. Vidyawati College of Pharmacy, Jhansi (U.P.) – 284121, India. Ocimum sanctum Linn: A Review on Phytopharmacology and Therapeutic Potential of Tulsi, International Journal of Pharmaceutical and Phytopharmacological Research (eIJPPR), Res. 2013; 3 (2): 148-151
- [4]. Mondal S., Bijay R. Miranda R. B., and Sushil C. M. The Science behind Sacredness of Tulsi (Ocimum sanctum LINN.). Ind J of Physiol Pharmacol. 2009, 53: 291–306.
- [5]. Naji-Tabasi S, Razavi SM. Functional properties and applications of basil seed gum: An overview. Food Hydrocolloids 2017.
- [6]. Muralikrishnan G, Pillai SK, Shakeel F. Protective effects of Ocimum sanctum on lipid peroxidation and antioxidant status in streptozocin-induced diabetic rats. Natural product research. 2012.
- [7]. Ahmad MZ, Ali M, Mir SR. Anti-diabetic activity of Ocimum sanctum L. roots and isolation of new phytoconstituents using two-dimensional nuclear magnetic resonance spectroscopy. J Pharmacogn Phytother. 2012;4:75–85. [Google Scholar]
- [8]. Singh PK, Baxi D, Banerjee S, Ramachandran AV. Therapy with methanolic extract of Pterocarpus marsupium Roxb and Ocimum sanctum Linn reverses dyslipidemia and oxidative stress in alloxan induced type I diabetic rat model. Exp Toxicol Pathol. 2012;64:441–8. [PubMed] [Google Scholar]
- [9]. Correa M.G., Couto J.S., Teodoro A.J. Anticancer properties of Psidium guajava—A mini-review. Asian Pac. J. Cancer Prev. 2016;17:4199–4204. [PubMed]
- [10]. Shu Y, Liu Y, Li L, Feng J, Lou B, Zhou X et al. Antibacterial activity of quercetin on oral infectious pathogens. Afr J Microbiol Res. 2011; 5:5358-61.10.
- [11]. Birdi T, Daswani P, Brijesh S, Tetali P, Natu A et al. Newer insights into the mechanism of action of Psidiumguajava L. leaves in infectious diarrhoea.
- [12]. BMCComplement Altern Med. 2010.
- [13]. Okoli CO, Ezike AC, Akah PA, Udegbunam SO, OkoyeTC, Mbanu TP, Ugwu E. Studies on wound healing &antiulcer activities of extract of aerial parts ofPhyllanthus niruri L.(Euphorbiaceae).Am J PharmacolToxicol. 2009.
- [14]. Lozoya X, Reyes-Morales H, Chavez-Soto M, Martínez-García Mdel C, Soto- González Y et al. Intestinal anti-spasmodic effect of a phytodrug of Psidium guajava folia in the treatment of acute diarrheic disease. JEthnopharmacol. 2002; 83:19- 24.26.
- [15]. Nisha, M. H. (2011). Phytochemical and Biological investigation of Tridax procumbens leaves. Thesis report submitted to the Department of Pharmacy, East West University, Bangladesh, in partial fulfillment of therequirements for the degree in B. Pharm.

- [16]. Policegoudra, R. S., Chattopadhyay, P., Aradhya, S. M., Shivaswamy, R., Sing, L., & Veer, V. (2014). Inhibitory effect of *Tridax procumbens* against human skin pathogens. *Journal of Herbal Medicine*, 4(2), 83-88. <https://doi.org/10.1016/j.hermed.2014.01>
- [17]. Manjamalai, A., & Grace, V. M. B. (2004). Effect of essential oil of *Tridax procumbens* Linn on in-vivo antioxidant level in cancer model and in-vitro free radical scavenging activity. *International Journal of Pharmaceutical Analysis*.
- [18]. Dhanavade MJ, Jalkute CB, Ghosh JS, Sonawane KD (2011) Study antimicrobial activity of lemon (*Citrus lemon* L.) peel extract. *Br J Pharmacol Toxicol* 2: 119-122. Link: <https://bit.ly/2ZoDkzX>
- [19]. Robards K., Antolovich M. Analytical chemistry of fruit bioflavonoids. A review. *Analyst*. 1997;122:11–34. doi: 10.1039/a606499j.
- [20]. Raimondo S., Naselli F., Fontana S., Monteleone F., Lo Dico A., Saieva L., Zito G., Flugy A., Manno M., Di Bella M.A., et al. Citrus limon-derived nanovesicles inhibit cancer cell proliferation and suppress CML xenograft growth by inducing TRAIL-mediated cell death. *Oncotarget*. 2015;6:19514–19527. doi: 10.18632/oncotarget.4004.
- [21]. Millet F. Huiles essentielles et essence de citronnier (*Citrus limon* (L.) Burm. f.) *Phytotherapie*. 2014;12:89–97. doi: 10.1007/s10298-014-0857-3. [CrossRef]
- [22]. Kato Y., Domoto T., Hiramitsu M., Katagiri T., Sato K., Miyake Y., Aoi S., Ishihara K., Ikeda H., Umei N., et al. Effect on blood pressure of daily lemon ingestion and walking. *J. Nutr. Metab.* 2014;2014 doi: 10.1155/2014/912684
- [23]. Journal of Novel Research in Pharmacy and Technology (JONRPT)
- [24]. Journal of Novel Research in Pharmacy and Technology (JONRPT)
- [25]. Okwu, D.E.; Ezenagu, V.I.T.U.S. Evaluation of the phytochemical composition of mango (*Mangifera indica* Linn) stem bark and leaves. *Int. J. Chem. Sci.* 2008, 6, 705–716.
- [26]. Zhu XM, Song JX, Huang ZZ, Whu YM, Yu MJ. Antiviral activity of mangiferin against herpes simplex virus type 2 in vitro. *Zhongguo Yao Li Xue Bao*. 1993;14:452–[PubMed] [Google Scholar]
- [27]. Zheng MS, Lu ZY. Antiviral effect of mangiferin and isomangiferin on herpes simplex virus. *Chin Med J*. 1990;103:160–5.
- [28]. Rivera DG, Balmaseda IH, Leon AA, Hernandez BC, Montiel LM, Garrido GG, et al. Anti-allergic properties of *Mangifera indica* L. extract (Vimang) and contribution of its glucosylxanthone mangiferin. *J Pharm Pharmacol.* 2006;58:385–92.
- [29]. Muanza DN, Euler KL, Williams L, Newman DJ. Screening for antitumor and anti-HIV activities of nine medicinal plants from Zaire. *Int J Pharmacol.* 1995;33:98.
- [30]. Tomar Lokeshwar et al. IRJP 2011, 2 (12) 97-102.
- [31]. Pingale Shirish S, Hepatoprotection study of leaves powder of *Azadirachtaindica*juss, International Journal of Pharmaceutical Sciences Review and Research, Volume 3, Issue 2, July – August. 2010.
- [32]. Rehman H, Yasin KA, Choudhary MA, Khaliq N, Rahman A, Choudhary MI, Malik S, Studies on the chemical constituents of *Phyllanthus emblica*, *Natural Product Research*, 21(9), 2007, 775-81.
- [33]. Krishnaveni M, Mirunalini S, Therapeutic potential of *Phyllanthus emblica* (amla): the ayurvedic wonder, *Journal of Basic and Clinical Physiology and Pharmacology*, 21, 2010, 93-105.
- [34]. Sankaran M, Velusamy V, Mani K, Amla: a novel ayurvedic herb as a functional food for health benefits”- A mini review, *International Journal of Pharmacy and Pharmaceutical Sciences*, 5(1), 2013, 1-4.
- [35]. Verma V, Bakshi SK. *Tinospora cordifolia* Miers ex Hook.f & Thoms. Medicinal and aromatic plants. http://portal.ics.trieste.it/MAPs/MedicinalPlants_Plant.aspx?id=660 [Accessed March 20, 2011].
- [36]. Asthana JG, Jain S, Mishra A, Vijaykant MS. Evaluation of antileprotic herbal drug combinations and their combination with Dapsone. *Indian Drugs* 2001;38:82-6.
- [37]. Zhao TF, Wang X, Rimando AM, Che C. Folkloric medicinal plants: *Tinospora sagittata* var. *cravaniana* and *Mahonia bealei*. *Planta Med* 1991;57:505.
- [38]. Pramila Ghumare, Jirekar DB, Mazahar Farooqui, Naikwade SD. A Review of *Pongamia pinnata* – An Important Medicinal Plant. *Current Research in Pharmaceutical Sciences*. 4 (02), 2014,

- [39]. Tripathi S, Kumar H, Pant H. Wood protection by Pongamia pinnata seed oil through heating process. Indian Forester. 138, 2012, 136–41.
- [40]. Li J, Jiang Z, Li X, Hou Y, Liu F, Li N, Liu X, Yang L, Chen G. Natural therapeutic agents for neurodegenerative diseases from a traditional herbal medicine Pongamia pinnata (L.) Pierre. Bioorganic & medicinal chemistry letters. 2015 Jan 1, 53-8.
- [41]. Atherton P. Aloe vera revisited. Br J Phytother. 1998;4:76–83. [Google Scholar]
- [42]. Shelton M. Aloe vera, its chemical and therapeutic properties. Int J Dermatol. 1991;30:679–83. [PubMed] [Google Scholar]
- [43]. Ishii Y, Tanizawa H, Takino Y. Studies of aloe. V: Mechanism of cathartic effect. Biol Pharm Bull. 1994;17:651–3. [PubMed]
- [44]. Kim HS, Lee BM. Inhibition of benzo [a] pyrene-DNA adduct formation by aloe barbadensis Miller. Carcinogenesis. 1997;18:771–6
- [45]. Tripathi and Ruchika Sharma, Bio-chemical activities of Ficus banghalensis- A review article.
- [46]. Mandal, S. G., Shete, R. V., Kore, K. J., Otari, K. V., Kale, B. N. and Mamma, A.K, 2010. Review: Indian national tree (Ficus bengalensis), International Journal of Pharmacy & Life Sciences, 1, (05), 268-273.
- [47]. Tripathi and Ruchika Sharma, Bio-chemical activities of Ficus banghalensis- A review article
- [48]. Tripathi and Ruchika Sharma, Bio-chemical activities of Ficus banghalensis- A review article
- [49]. Costa-Luatufo LV, Khan MTH, Ather A, WilkeDV, Jimenez PC, Pessoa C, Moaraes MEA and Moreas MOD, Studies of anticancer potential of plants used in Bangladeshi folk medicine,J Ethnopharmacol, 2005, 99(1) 21-30.
- [50]. Arul V, Miyazaiki S and Dahananjaya R, Studieson the anti-inflammatory, antipyretic,analgesic properties of the leaves of Aegle marmelos, J Ethnopharmacol, 2005,
- [51]. Paricha S, Bael (Aegle marmelos) Nature'smost natural Medicinal Fruit, Orissa Rev,2004, 16-17.
- [52]. Roy SK and Singh RN, Studies on changesduring development and ripening of Bael fruit, Punjab Hortic J, 1980, 20(3&4),190-197.
- [53]. Indian journal of applied research.623 .
- [54]. Charde R.M, Dhongade H.J, Charde M.S, Kasture A.V. (2010).Evaluation of antioxidant, wound healing and anti-inflammatory activity of ethanolic extract of leaves of Ficus religiosa. International Journal of Pharma Sciencesand Research, 1: 73-82.
- [55]. Kaur H, Singh D, Singh B, Goel R.K. (2010). Anti-amnesic effect of Ficus religiosa in scopolamine-induced anterograde and retrogradeamnesia. Pharm Biol, 48, 234–240. 46. Kaushik R.K,
- [56]. de Amorim A, Borba H.R, Carauta J.P, Lopes D, Kaplan M.A. (1999). Anthelmintic activity of the latex of Ficus species.J Ethnopharmacol, 64:255–8. 48. Mallurvar V & Pathak A.K. (2008). Studies on immunomodulatory activity of Ficus religiosa. Indian
- [57]. Ganesan P, Phaiphan A, Murugan Y, Baharin B.S., Comparative study of bioactive compounds in curryand coriander leaves: An update, Journal of Chemical and Pharmaceutical Research 2013.
- [58]. Parul S, Javed A, Neha B, Honey J, Anuj B., Curry leaves- A medicinal herb, Asian Journal ofPharmaceutical Research., 2012, 2(2),
- [59]. Jean Bruneton, Carica papaya, In:Pharmacognosy, Phytochemistry of Medicinalplants, 2nd Edn, Technique & Documentation,
- [60]. Bhat Praveen G and Surolia Namita, In vitro Antimalarial activity of extracts of three plants used in the traditional medicine of India, AmJ Trop Med Hyg, 2001, 65(4), 304-308
- [61]. Calzada F, Yepez-Mulia L and Tapia-Contreras A, Effect of Mexican medicinal plant used to treat trichomoniasis on Trichomonas vaginalis trophozoites,J Ethnopharmacol, 2007, 113(2)
- [62]. tenuiflorum (tulsi) is an aromatic plant in the family Lamiaceae