

# **Productivity and Proximate Composition of *Pleurotus flabellatus* Cultivated on Different Agro-Wastes**

**Yadav S. G.**

Department of Botany

Shivaji Mahavidyalaya, Renapur, Latur, Maharashtra, India

**Abstract:** *Pleurotus flabellatus* was cultivated on selected agro wastes viz. soybean, paddy, green gram straw and their combinations in 1:1 proportion to study the productivity and nutritional content (moisture, crude protein, total carbohydrates, fat, crude fiber, ash and minerals like Ca, P, Fe.) of mushroom. Soybean straw, paddy straw and combination of soybean + paddy straw were found to be more suitable than green gram straw. The bioconversion of different agro wastes by *P. flabellatus* results into an improved human food.

**Keywords:** Proximate composition, *P. flabellatus*, Bio-efficiency, agro wastes

## **REFERENCES**

- [1]. Akindahunsi, A. A. and Oyetayo, F. L. (2006). Nutrient and antinutrient distribution of edible mushroom, *Pleurotus tuber-regium* (Fries) Singer. LWT,39,548-553
- [2]. Anonymous (1980) A Practical Manual of Nutrition, NIN. National Institute of Nutrition, Hyderabad
- [3]. AOAC (1990). Official methods of analysis (15th ed.) .Washington, DC : Association of Official Analytical Chemists.
- [4]. Bano, Z. and Srivastava, H.C. (1962) Studies in the cultivation of *Pleurotus* Sp. on Paddy straw. Food Sci. 12 : 362 - 368
- [5]. Buigut, (2002). Musroom production in sustainable small scale farming system opportunities and constraints : a survey of Uasin Gishu district. In : Proceedings of the Horticulture seminar on Sustainable Horticultural production in the Tropics at Jomo Kenyatta University of Agri. & Technologi, Juja, Kenya 3rd - 6th October, 2001 Eds, Wesonga JM, Losenge T, Ndungu's CK, Fricke, A. Han, B., Stutz, H. pp-1-5
- [6]. Chang, S.T., Lau, O.W. and Cho, K. Y. (1981). The cultivation and nutritive value of *Pleurotus sajor-caju*. European J. Appl. Microbiol. Bio technol. 12 : 58-62.
- [7]. Mandhare, V.K. (2000). Productivity of *Pleurotus* sp. on different substrates and its effect on Nutritional Indices of Spent straw. Ph.D. Thesis. Marathwada Agriculture University, Parbhani.
- [8]. El-Kattan, M.H.; Helmy, Z.A.; Abdel, H.; El-Liethy, M. and Abdelkawi, K. A. (1991) Studies on cultivation techniques and chemical composition of oyster mushrooms. Mushroom J. for tropics. 11 (3/4): 59-66.
- [9]. Kadlag, G.K.; Wani, P.V. and Sawant, D.M. (1998) Comparative performance of different *Pleurotus* species on wheat and green gram straw. J. Maharashtra Agric. Univ. 23 (1): 85-86.
- [10]. Necla, Caglarirmak (2007). The nutrients of exotic Mushrooms (*Lentinus edodes*, *Pleurotus* spp.) and an estimated approach to the volatile compounds. Food Chemistry. Article in Press.
- [11]. Patil, S.S. and Dakore,H.G. (2007). Comparative study on yield performance and Nutritive value of Oyster mushroom on Soybean straw. Bioinfolet 4 (1) : 57-59.
- [12]. Panse, V.G. and Sukhatme, P.V. (1978). Stat. Math. Agri. Work. ICAR publication, New Delhi.
- [13]. Ranganna, S. (1986) handbook of analysis and Quality Control for fruit and vegetable products, Tata McGraw Hill Publ. Company Ltd., New Delhi.
- [14]. Wankhede, D.B. and Tharanathan, R. N. (1976). Sesame (*Sesamum indicum*) Carbohydrates. J. Agri. Food. chem. 24 : 655 – 659.