

Review on Rauwolfiaserpentina Mother Tinture

Mundhe Siddhika S¹, Kokane Snehal M², Inamdar Saniya N³, Shinde Pallavi M⁴, Datkhile Sachin V⁵

Students, Samarth Institute of Pharmacy, Belhe, Maharashtra, India^{1,2,3}

Department of Pharmaceutics, Samarth Institute of Pharmacy, Belhe, Maharashtra, India^{2,5}

mundhesiddhika@gmail.com

Abstract: Root of sarpagandha could be a species of seed plant within the dogbane family has been traditionally employed in Ayurveda for many years to treat the variability of diseases of that at been traditionally apper in reality little the same as each other. rauvolfia is a safe and effective treatment of hypertension.much smaller dose of reserpine is required to get the antihypertensive activity.the present review focuses mainly on chemical composition, pharmacology, mechanism of action,side effects and toxicity and antihypertensive effect of Rauwolfia alkaloids. The plant provide clinician with safe and effective adjunct to high vital sign.

Keywords: Rauwolfia, Hypertension, Reserpine

REFERENCES

- [1]. Mollar J. M. schittler E, et al, “ reserpine, Der sedative Wirkstoffrauwolfia serpentine (benth) experientia” 1952;8: 338.
- [2]. Deshmukh S.R. Dhanshree SA, patil BA, Extraction and Evaluation of indole alkaloids from rauwolfiaserpentina For their antimicrobial and antiproliferative activities, International journal of pharmacy and pharmaceutical Sciences, 4(5), 2012, 329-334.
- [3]. Tyler V.E.,Brady L.R., Robbers J.E.,Pharmacognosy, 9th edition, Lea and Febiger,Philadelphia,1988,222-224
- [4]. Vakil R.J., “Rauwolfiaserpentina in the treatment ofHigh Blood Pressure: Review of the Literature”, Circulation, 1955, 12:220-229.
- [5]. Endress ME, Bruyns PV. A revised classification of the Apocynaceae. I. Bot Rev. 2000;66(1):1–56.
- [6]. E.G. Mc Queen, A.E. Doyle, et al, “Mechanism of Hypotensive Action of Reserpine, an alkaloid of Rauwolfiaserpentina”, Nature, 1954,174, 1015.
- [7]. Schuldiner S, Liu Y, Edwards RH. Reserpine binding to A vesicular amine transporter expressed in ChineseHamster ovary fibroblasts. J Biol Chem. 1993;268(1):29–34.
- [8]. Wimalasena K. Vesicular monoamine transporters: Structure-function, pharmacology, and medicinalChemistry. Med Res Rev.2011;31(4):483–519.
- [9]. Panwar GS, Guru SK. Alkaloid profiling and estimation Of reserpine in Rauwolfiaserpentina plant by TLC, HP-TLC and HPLC. Asian J Plant Sci. 2011;10(8):393–400.
- [10]. Jerie P. Milestones of cardiovascular therapy, IV: Reserpine [in Czech] CasLekCesk. 2007;146(7):573–577.
- [11]. Bein, H.J. “ZurPharmakologie des Reserpine, einesNeuenAlk.ausRauwolfiaserpentina (Benth)”, Experientia, 1953, 9: 1073.
- [12]. Goto, Y., Personal Communication from Tokyo, Japan.
- [13]. VakilR.,“Hypotensive action of other RauwolfiaSerpentina alkaloids: a clinical report”, J. Indian M.A., 1953, 23: 97-102.
- [14]. Guy Lemieux, M.D., Andre Davignon, M.D., et al,” Treatment of arterial hypertension with Rescinnamine, A new alkaloid isolated from Rauwolfia serpentina”,956, 74, 145.
- [15]. Yarnell E, Abascal K. Treating hypertension Botanically. Altern Complement Ther. 2001;7(5):284–290.
- [16]. Krosgaard AR. The effect of reserpine on the Electrolyte and fluid balance in man. Acta Med Scand. 1957;159(2):127–132.
- [17]. Baumeister AA, Hawkins MF, Uzelac SM. The myth of Reserpine-induced depression: role in the historical Development of the monoamine hypothesis. J HistNeurosci. 2003;12(2):207–220.

