IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 1, December 2022

Review On: Lyophilization Process of Pharmaceuticals Manufacturing

Pranali Kulawade, Aditi Lende, Shweta Mule, Rutuja Bhand, Mrunal Mahajan, Gaytri Korde, Sanket Jawale, Minaj Inamdar, Swaranjali Shinde

Department of Pharmaceuticas, Samarth College of Pharmacy, Belhe, Pune, Maharashtra, India kulawadepranali@gmail.com

Abstract: To establish a commertially feasible shelf life, lyophillization (freeze drying) is frequently employed to manufacture dry pharmaceuticals formulations. It is not necessary to refrigerate properly frozen dried products. Also kept at room temperature. Lyophilization is common, but it is expensive intense. Using the right process parameters, we can get the results we want. Compared to standard drying methods, our products are of the highest quality methods. Lyophilization has become increasingly significant in the pharmaceutical industry. It is subject to further growth and expansion. The procedure includes the following steps: Freezing, primary drying, and secondary drying are the three steps. Article focused on how various elements influence the lyophilization process, as well as provides a complete list of excipients used in lyophilized products...

Keywords: Lyophilization, Freezing, Primary, and Secondary Drying, End point of freeze drying

REFERENCES

- [1]. Kundra T "Mujumdar AS ., Advanced drying Technologies. 2nd ed. CRC Press; 2009.
- [2]. Lippincolt, Williams K. Remington, The Science & practice of pharmacy, Parenteral Preparation, 20th ed, ISE Publication phelabelohia .2000;1:804-819.
- [3]. Chien & Yiew W. Pharmaceutical Dosage forms: Parenteral Medications. Indian Journal of pharmaceutical science and technology, 1981; 35: 106-118.
- [4]. Oetjen GW., Freeze-Drying, Encycl. Sep. Sci. Elsevier Science 2000; 1023–34.
- [5]. Genin N., Rene F., Corrieu G., A method for on-line Determination of residual water content and sublimation End-point during freeze-drying, Chem. Eng. Process. Process Intensif. 1996; 35: 255–63.
- [6]. Charles P, Detke HC, Pyne A. Post injection delirium/sedation syndrome in patients with Schizophrenia treated with Olanzapine long acting injection: analysis of cases. BMC Psychiatry, 2005.
- [7]. Nail SL, Johnson W, Methodology for in-process determination of residual water in Freezedried products. Dev Biol Stand.1992; 74: 137–51.
- [8]. Roy M., Pikal MJ, Process control in freeze drying: determination of the end point of Sublimation drying by an electronic moisture sensor. J Parenter Sci Technol. 1989; 43(2): 60.
- [9]. Nail SL, Gatlin GA. Freeze drying: principles and practice. Marcel Dekker publisher, Newyork. 1992; 2: 163–233.
- [10]. Dagleish MJ & Swarbrick J. Encyclopedia of Pharmaceutical Technology Volume 3,Informa Healthcare publisher, USA. 2007; 1807-1833.
- [11]. Remington: The science and practice of pharmacy, 21st ed, Gennaro RA, Lippincott c illiams & wilkins publisher, 2000; 1.
- [12]. Jeff SJ. Basic Cycle Development Techniques for Lyophilized Products. 2009; 35: 126-128.
- [13]. Adams GD, Irons LI. Some implications of structural collapse during freeze drying using Erwinia caratovora l-asparaginase as a model. J Chem Biotechnol, 1993; 58: 71–76.
- [14]. Sanjith NL & Gatin LA. Freeze drying: Annealing principles and practice. NP publication. 1993; 2: 163-233.

IJARSCT



International Journal of Advanced Research in Science, Communication and Technology (IJARSCT)

Volume 2, Issue 1, December 2022

- [15]. Neema S., Washkuhn RJ and Brendel RJ; Excipients and their use in Injectable products; PDA J Pharm Sci Technol; 1997; 51; 166-171
- [16]. Shah N.V., Solanki H and Prajapati V; Impact of formulation ingredients on quality of the parenteral products; WJPPS; 2015; 4(3); 468-482
- [17]. Shah R and Mehta P; Freeze dried injectable drug product development selection of non-functional additives; Int J Pharm Pharm Sci; 2014; 6(9): 3-7
- [18]. Shireesh P.A and Sydney O.U; A review and classification of emerging excipients in parenteral medications; Pharm Tech; 2003; 46-60
- [19]. Sougata P., Deepak S and Vikas C; Excipient selection in parenteral formulation development; Pharma Times; 2013; 45(3); 65-77
- [20]. Yasir M and Umer F; Excipients use in parenteral and lyophilized formulation development; Open science journal of pharmacy and pharmacology; 2015; 3(3); 19-27
- [21]. Ankit B., Lokesh K and Arvind K.B; Excipients used in lyophilization of small molecules; J. Excipients and Food Chem; 2010; 1(1); 41-54

DOI: 10.48175/568