

International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 5, Issue 1, June 2025



# Formulation and Evaluation of Herbal Aloe Vera Gel

Miss. Pooja Meghraj Gunthal, Miss. Vaishnavi Shahaji Phatak, Miss. Geeta Narayan Tikule Dr. Surwase K. P.

Aditya Institute of Pharmaceutical College, Beed

Abstract: Aloe Vera, commonly known as Barbados or Curacao Aloe, is an herbal medicine with along tradition of use by a variety of cultures. The succulent plant grows in arid and subtropical climates and is best known for 2 distinct preparations: the clear mucilaginous gel that is widely used for the treatment of minor burns, especially sunburns, and the thick sap of the leaves that turns yellow-brown and has strong laxative effects that caution its use. The traditional uses of the clear mucilaginous gel are manifold, ranging from topical applications to reduce perspiration to oral dosing for diabetes and a range of gastrointestinal ailments. The efficacy of aloe Vera gel to treat burn wounds, genital herpes, and seborrhea dermatitis have been shown in clinical trials, but other indications such as psoriasis or internal application for the treatment of type 2 diabetes remain inconclusive. The main limitation of the current clinical trials are being conducted tofurther evaluate the use of aloe vera gel for a variety of disorders, as well as to further confirm traditional uses of the plant extract.

Keywords: Aloe vera, Barbados, dermatology

### I. INTRODUCTION

Aloe vera (syn. Aloe barbadensis Mill., Fam. Liliaceae), also known as Barbados or Curaçao Aloe, has been used in traditional and folk medicines for thousands of years to treat and cure a variety of diseases. Although the plant is native to northern parts of Africa, it has rapidly spread across the world because its cultivation is easy. An important distinction has to be made between the strongly laxative and purgative latex derived from the bundle-sheath cells and the clear mucilaginous gel. The plant has been used by Egyptians, Assyrians, and Mediterranean civilizations, as well as in Biblical times. A variety of aloe species are still used in folk medicines of Africa and Asia. Hunters in the Congo reportedly rub their bodies in the clear mucilaginous gel to reduce perspiration; some African tribes apply the gel for chronic conjunctivitis; the gel is used in India for the treatment of asthma.

Aloe vera gel is used as an ethnomedicine in Trinidad and Tobago for hypertension. The most common folk use of aloe has been for the treatment of burn wounds and specifically to aid in the healing process, reduce inflammation, and tissue scaring. The gel was described by Dioscorides and used to treat wounds and mouth infections, soothe itching, and cure sores. The use of aloe vera gel as a household remedy in the United States was triggered by reports of its beneficial effect on radiation dermatitis followed by a boom in cultivation in the 1930s; it remains a common plant and for burns and abrasions. Important contemporary uses of the gel exist in traditional medicines of India, China, and Mexico, as well as Middle America and the West Indies. Mexico is producing roughly 47% of aloe worldwide with a total sales volume of \$123.5 million US dollars as of 2008.

Despite its widespread popularity, scientific evidence on the aloe vera gel remains sparse. Aloe vera gel is regarded as safe if applied topical with only a few allergic reactions being reported. The efficacy of aloe vera gel to treat burn wounds, genital herpes, and seborrheic dermatitis have been shown in clinical trials, but other indications such as psoriasis or internal application for the treatment of type 2 diabetes remain inconclusive. The major application of aloe vera gel remains as a skin moisturizer in cosmetics and as an après treatment for sunburns, for which it has proven its effectiveness.

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



The Aloe vera plant has been known and used for centuries for its health, beauty, medicinal and skin care properties. The name Aloe vera derives from the Arabic word "Alloeh" meaning "shining bitter substance," while "vera" in Latin means "true." 2000 years ago, the Greek scientists regarded Aloe vera as the universal panacea. The Egyptians called Aloe "the plant of immortality." Today, the Aloe vera plant has been used for various purposes in dermatology.

### HISTORY

Aloe vera has been used for medicinal purposes in several cultures for millennia: Greece, Egypt, India, Mexico, Japan and China. Egyptian queens Nefertiti and Cleopatra used it as part of their regular beauty regimes. Alexander the Great, and Christopher Columbus used it to treat soldiers' wounds. The first reference to Aloe vera in English was a translation by John Goodyew in A. D. 1655 of Dioscorides' Medical treatise De Materia Medica. By the early 1800s, Aloe vera was in use as a laxative in the United States, but in the mid-1930s, a turning point occurred when it was successfully used to treat chronic and severe radiation dermatitis.

Aloe vera is a succulent plant with thick, fleshy, serrated, lanceolate- shaped leaves of green-greyish color. Aloe vera inner gel is obtained from the lower leaves of the plant by slicing the leaf open. The gel is clear, odorless, and tasteless and should be free of leaf skin or yellow parts. No consistent standardization has been established, but the International Aloe Science Council (IASC), a trade association of internationally based aloe producers and marketers, requires adherence to certain specifications for the product to be certified.10 Other preparations include a hydrophilic cream containing 0.5% aloe vera gel and an emulsion consisting of 30% aloe vera gel.

The botanical name of Aloe vera is Aloe barbadensis miller. It belongs to Asphodelaceae (Liliaceae) family, and is a shrubby or arborescent, perennial, xerophytic, succulent, pea- green color plant. It grows mainly in the dry regions of Africa, Asia, Europe and America. In India, it is found in Rajasthan, Andhra Pradesh, Gujarat, Maharashtra and Tamil Nadu.

The plant has triangular, fleshy leaves with serrated edges, yellow tubular flowers and fruits that contain numerous seeds. Each leaf is composed of three layers:

1) An inner clear gel that contains 99% water and rest is made of glucomannans, amino acids, lipids, sterols and vitamins

2) The middle layer of latex which is the bitter yellow sap and contains anthraquinones and The outer thick layer of 15–20 cells called as rind which has protective function and synthesizes carbohydrates and proteins. Inside the rind are vascular bundles responsible for transportation of substances such as water (xylem) and starch

### **MECHANISM OF ACTIONS**

1. Healing properties: Glucomannan, a mannose-rich polysaccharide, and gibberellin, a growth hormone, interacts with growth factor receptors on the fibroblast, thereby stimulating its activity and proliferation, which in turn significantly increases collagen synthesis after topical and oral Aloe vera. Aloe gel not only increased collagen content of the wound but also changed collagen composition (more type III) and increased the degree of collagen cross linking. Due to this, it accelerated wound contraction and increased the breaking strength of resulting scar tissue. An increased synthesis of hyaluronic acid and dermatan sulfate in the granulation tissue of a healing wound following oral or topical treatment has been reported.

2. Effects on skin exposure to UV and gamma radiation: Aloe vera gel has been reported to have a protective effect against radiation damage to the skin. Exact role is not known, but following the administration of aloe vera gel, an antioxidant protein, metallothionein, is generated in the skin, which scavenges hydroxyl radicals and prevents suppression of superoxide dismutase and glutathione peroxidase in the skin. It reduces the production and release of skin keratinocyte-derived immunosuppressive cytokines such as interleukin-10 (IL-10) and hence prevents UV-induced suppression of delayed type hypersensitivity.

3. Anti-inflammatory action: Aloe vera inhibits the cyclooxygenase pathway and reduces prostaglandin E2 production from arachidonic acid. Recently, the novel anti- inflammatory compound called C-glucosyl chromone was isolated from gel extracts.

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



4. Effects on the immune system: Alprogen inhibit calcium influx into mast cells, thereby inhibiting the antigenantibody-mediated release of histamine and leukotriene from mast cells. In a study on mice that had previously been implanted with murine sarcoma cells, acemannan stimulates the synthesis and release of interleukin-1 (IL-1) and tumor necrosis factor from macrophages in mice, which in turn initiated an immune attack that resulted in necrosis and regression of the cancerous cells. Several low-molecular-weight compounds are also capable of inhibiting the release of reactive oxygen free radicals from activated human neutrophils.

5. Laxative effects: Anthraquinones present in latex are a potent laxative. It increases intestinal water content, stimulates mucus secretion and increases intestinal peristalsis.

6. Antiviral and antitumor activity: These actions may be due to indirect or direct effects. Indirect effect is due to stimulation of the immune system and direct effect is due to anthraquinones. The anthraquinone aloin inactivates various enveloped viruses such as herpes simplex, varicella zoster and influenza. In recent studies, a polysaccharide fraction has shown to inhibit the binding of benzopyrene to primary rat hepatocytes, thereby preventing the formation of potentially cancer-initiating benzopyrene-DNA adducts. An induction of glutathione S-transferase and an inhibition of the tumor- promoting effects of phorbol myristic acetate has also been reported which suggest a possible benefit of using aloe gel in cancer chemoprevention.

7. Moisturizing and anti-aging effect: Mucopolysaccharides help in binding moisture into the skin. Aloe stimulates fibroblast which produces the collagen and elastin fibers making the skin more elastic and less wrinkled. It also has cohesive effects on the superficial flaking epidermal cells by sticking them together, which softens the skin. The amino acids also soften hardened skin cells and zinc acts as an astringent to tighten pores. Its moisturizing effects has also been studied in treatment of dry skin associated with occupational exposure where aloe vera gel gloves improved the skin integrity, decreases appearance of fine wrinkle and decreases erythema. It also has anti-acne effect.

8. Antiseptic effect: Aloe vera contains 6 antiseptic agents: Lupeol, salicylic acid, urea nitrogen, cinnamonic acid, phenols and sulfur. They all have inhibitory action on fungi, bacteria and viruses.



Fig.1 Alovera plant leaves

### LITERATURE REVIEW

1) Aggarwal D and Barna K. (2004).

Micropropagation protocol for an elite selection of Aloe vera syn A. barbadensis through enhanced axillary branching was standardized. Murashige and Skoog medium containing 1 mg l–1 BA and 0.2 mg l-1 IBA gave highest multiplication. Citric acid at 10mg l-1 and liquid medium improved the shoot multiplication. Hundred per cent microshoots produced rooted plantlets within 15 days of culture on hormone-free agar medium. Liquid medium during rooting stage decreased the number of shoots showing rooting response. The plants were successfully transferred in the soil and were morphologically similar to mother plants

#### 2) Barna, K.S., Wakhlu, A.K. 1994.

A protocol for whole plant regeneration of Cicer arietinum L. cv. C-235 via organogenesis from callus has been developed. Callus initiation was best when immature leaflets were cultured on MS medium containing 5 or 25  $\mu$ M 2,4-

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



D or NAA in combination with 10  $\mu$ M BA, or 25  $\mu$ M 2,4-D alone. The callus grew most vigorously on MS medum supplemented with 10  $\mu$ M NAA and 5  $\mu$ MBA. Best shoot differentiation was obtained from calli derived from the basal portion of shoot tips on MS medium supplemented with 10  $\mu$ M BA and 0.1  $\mu$ M IBA. The shoot forming ability of calli was enhanced by adding 5 mM potassium phosphate to the medium. Shoots were rooted on a MS medium containing 1  $\mu$ M IBA. The regenerated plants were grown to maturity and produced viable seed.

#### 3) S.AHMED, A.H. KABIR, M.B. AHMEDI, M.A. RAZVY2 and S. GANESAN [2007]

method has been developed using shoot tip explants in Aloe vera L. The process involves subsequent in vitro morphogenesis and rooting of the in vitro proliferated shoots and transplantation of regenerated plants under ex vitro condition. Shoot proliferation was found best in MS medium containing BA 2.0 mg/l, KIN 0.5 mg/l and NAA 0.2 mg/l. Maximum 98.96% shoots were proliferated in this media composition. This media composition is best comparing to other treatment used in this study.

#### 4) N. Dwivedi, A. Indiradevi, K. Asha, N. Asokan, and A. Suma,

The research article by Dwivedi et al. outlines a protocol for micropropagation of Aloe vera (Indian Aloe) using axillary shoots. It demonstrates that these shoots, when cultured in MS medium supplemented with 1.5 mg/l BAP, produce new plants and roots simultaneously. Each explant generates an average of 14 shoots and roots within 8 weeks, and this multiplication can continue for at least a year. The regenerated plantlets are successfully transferred to soil with an 83% survival rate.

#### 5) Steenkamp V and Stewart MJ. (2007).

The 2007 study by Steenkamp and Stewart, published in Pharmaceutical Biology, reviews the diverse med icinal applications and toxicological activities of Aloe products, specifically focusing on Aloe vera and Aloe ferox. The study highlights that Aloe, particularly its gel and latex, has been used for centuries for various purposes, including as a laxative, in topical treatments for skin conditions, and for treating a range of diseases.

#### 6) Chand, S. and Roy, S.C., 1981

In their 1981 study in Annals of Botany, Chand and Roy successfully induced organogenesis in callus cultures of Nigella sativa L. (black cumin). This means they were able to guide the growth of callus cells to form new plant organs like shoots and roots. Their research demonstrated the potential of Nigella sativa for in vitro plant regeneration through callus cultures.

#### 7) Durzan, D.J., 1984.

No part of this work may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, microfilming, recording or otherwise, without written permission from the Publisher, with the exception of any material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work.

#### 8) Schaik AH, Van Struik PC, Damian TG

Two experiments were carried out to investigate the effects of irrigation and N application on the initiation and growth of leaves, and thus gel production, of Aloe barbadensis Mill. under the typical arid conditions of Aruba. Irrigation proved to be essential for continuous vegetative growth and high yields of A. barbadensis in Aruba throughout the year. It increased the fresh weight per leaf, but, during the first months of drought, water deficiency hardly affected the leaf appearance rate. Vegetative reproduction by suckers was reduced by drought.

### 9) Natali L, Sanchez IC, Cavallini A

**Copyright to IJARSCT** 

www.ijarsct.co.in

The provided reference, "Natali L, Sanchez IC, Cavallini A. In vitro culture of Aloe barbadensis Miller micropropagation from vegetative meristems. Plant Cell Tiss. Org. 1990;20:71-74," describes a study on the in vitro

DOI: 10.48175/IJARSCT-27366







International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



micropropagation of Aloe barbadensis (Aloe vera) using vegetative meristems. This research likely involved culturing shoot tips or meristems in a controlled laboratory environment to produce multiple new plants, which is a common technique for mass-producing plants, according to scispace.com.

10) Meyer HJ, Staden

The text describes a scientific paper by Meyer HJ and Staden JV, published in 1991. The paper focuses on rapid in vitro propagation of Aloe barbadensis Miller, commonly known as aloe vera. It appeared in the journal "Plant Cell

# AIM AND OBJECTIVE

Aim:

A Brief Review on Formulation and Evaluation of Aloe Vera Gel by Using Natural Ingredient.

Objective of the study:

1. To achieve sustained improvement in Aloe-Vera gel compliance ratio.

2. To kill germs and microorganism that can harm our body.

3. It will help to learn their illness are often caused by germs which travels from their hands to their mouth, eye, nose etc.

Reduces the rates of healthcare associated infections

# **DRUG PROFILE**

Aloe vera contains 75 potentially active constituents:

Vitamins, enzymes, minerals, sugars, lignin, saponins, salicylic acids andamino acids.

1. Vitamins: It contains vitamins A (beta-carotene), C and E, which are antioxidants. It also contains vitamin B12, folic acid, and choline. Antioxidant neutralizes free radicals.

2. Enzymes:

- It contains 8 enzymes:
- Aliiase
- Alkaline phosphatase
- Amylase
- Bradykinase
- Carboxypeptidase
- Catalase
- Cellulose
- Lipase.

Bradykinase helps to reduce excessive inflammation when applied to the skin topically, while others help in the breakdown of sugars and fats.

3. Minerals: It provides calcium, chromium, copper, selenium, magnesium, manganese, potassium, sodium and zinc. They are essential for the proper functioning of various enzyme systems in different metabolic pathways and few are antioxidants.

4. Sugars: It provides monosaccharides (glucose and fructose) and polysaccharides: (glucomannans/polymannose). These are derived from the mucilage layer of the plant and are known as mucopolysaccharides.

The most prominent monosaccharide is mannose-6-phosphate, and the most common polysaccharides are called glucomannans. Acemannan, a prominent glucomannan has also been found. Recently, a glycoprotein with antiallergic properties, called alprogen and novel anti-inflammatory compound, C-glucosyl chromone, has been isolated from Aloe vera gel.



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



5. Anthraquinones: It provides 12 anthraquinones, which are phenolic compounds traditionally known as laxatives. Aloin and emodin act as analgesics, antibacterials and antivirals.

6. Fatty acids: It provides 4 plant steroids; cholesterol, campesterol,  $\beta$ - sisosterol and lupeol. All these have antiinflammatory action and lupeol also possesses antiseptic and analgesic properties.

7. Hormones: Auxins and gibberellins that help in wound healing and have anti- inflammatory action.

8. Others: It provides 20 of the 22 human required amino acids and 7 of the 8 essential amino acids. It also contains salicylic acid that possesses anti- inflammatory and antibacterial properties. Lignin, an inert substance, when included in topical preparations, enhances penetrative effect of the other ingredients into the skin. Saponins that are the soapy substances form about 3% of the gel and have cleansing and antiseptic properties.

### PLAN OF WORK :-



### MATERIAL AND METHOD

Material:

In Aloe vera powder, the carbohydrate content was determined according to mannose and phenolic compounds in terms of gallic acid.

- Aloe vera pulp
- Liquid glucose
- Glycerin

Weight uniformity, content uniformity, the organoleptic properties evaluation, releasing the active ingredient in the phosphate buffer (pH, 6.8) and taste evaluation were examined by Latin square method.

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



Plant Material

Aloe-vera, was taken as plant material in this formulation. Aloe-vera was collected from the Botanical Garden, Aditya Institute of Pharmaceutical, Beed.

Aloe-Vera

Aloe vera is a popular medicinal plant with antioxidant and antibacterial properties. It may reduce dental plaque, speed wound healing, prevent wrinkles, and manage blood sugar.

Benefits of Aloe-Vera

- Soothes Burns
- Heals Wounds
- Eases Intestinal Problems
- Reduces Arthritic Swelling
- Heals Psoriasis Lesions [19]
- Gum Infections
- Eye irritations and injuries
- Strains and sprains

### Chemicals

1] Vitamin E and Glycerin was collected from the chemical store room, Aditya Institute of Pharmaceutical, Beed.

2] Xanthan gum:-

xanthan gum is used as a thickening and stabilizing agent in foods, toothpastes, and medicines.

3] Glycerin :-

Glycerine helps reduce itching of the scalp fast. Aloe vera offers relief to skin and hair damaged by sunlight exposure. With organic Aloe vera, the skin spots and rashes fade away gradually.

Aloe vera locks moisture, whereas Glycerine brings radiance to the skin and hair.

4] Agar agar

Commonly employed in desserts like jellies, puddings, and custards, as well as in savory dishes to thicken soups, sauces, and even pie fillings.

5] Rose water

Calming skin irritation, reducing redness, and even treating mild infections

# Method:

• A few Aloe vera leaves from an Aloe vera (Barbadensis Miller) plant were collected and washed with water thoroughly.

• A sharp knife was taken and carefully peeled off the yellow layer just beneath the green rind avoiding the vascular bundles and the top rind was removed.

• The bottom rind was similarly removed to discard the significant amount of mucilage which gets attached onto this.

• The clear mucilaginous gel was scooped out with a spoon. 50 ml crude clear Aloe vera extract was mixed uniformly by using magnetic stirrer and the plant extract was filtered through a whatman filter paper.

• Finally, it was transferred into a sterilized, clean glass jar and stored in the refrigerator



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 5, Issue 1, June 2025









2. Aloe-Vera Leaves



3. Leaf under Cold water

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 5, Issue 1, June 2025





4. Slice off the outer layer



5. Scooping the Gel



6. Placing the Gel into Mortar to homogenize it

### FORMULATION OF ALOE VERA GEL

• The gel base was prepared by dispersing 1% xanthan gum in distilled water at 80°C with constant stirring at a moderate speed using magnetic stirrer and pH was adjusted to 6-7 using triethanolamine.

DOI: 10.48175/IJARSCT-27366

- 50 ml of Aloe vera extract was added into it and gel base using Aloe vera extract was prepared.
- The oil phase of the emulsion was prepared by dissolving 0.5 ml span 20 in 7.5 ml coconut oil.
- 1% of zinc oxide was added into it. Zinc oxide protects skin from UV induced damage.

Copyright to IJARSCT www.ijarsct.co.in







International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



- The aqueous phase was prepared by dissolving 1 ml tween 20 in purified water.
- Required amount of methyl paraben was added into aqueous phase and propyl paraben was dissolved in oil phase.
- Two phases were separately heated to 70-80 °C.

• The oil phase was added drop wise to aqueous phase and was mixed together with continuous stirring using mechanical stirrer at specified speed until it cooled down to room temperature. The prepared emulsion was mixed with the gel in 1:1 ratio with gentle stirring to obtain the Aloe vera gel.



Prepared Aloe Vera Gel

# ALOE VERA USES

External

- Mild to moderate burns as well as erythema
- Genital herpes
- Seborrheic dermatitis

### Internal:

• Adjunct therapy of spontaneous fibrosarcomas in dogs and cats.

Other Potential Uses:

(Determined by clinical trials and/or official monographs and/or empirical use)

- Psoriasis vulgaris
- Skin moisturizer
- Type 2 diabetes
- · Malignancies and immunodeficiency viruses in cats
- Oral lichen planus infections
- Angina pectoris
- Ulcerative colitis
- UV-induced erythema14 Kidney stones
- Alveolar osteitis

Clinical uses:

The clinical use of aloe vera is supported mostly by anecdotal data. Though most of these uses are interesting, controlled trials are essential to determine its effectiveness in all the following diseases

Uses based on scientific evidence Uses based on tradition or theory

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366







International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

Volume 5, Issue 1, June 2025







Fig.9 uses of aloevera

### **RESULTS AND DISCUSSION**

The aloe vera gel was evaluated as follows:

(i) Percentage Moisture Content:

Percentage moisture loss from the formulations were determined by the method reported by Deviet al. Two gram formulations were weighed (A1, A2, A3and A4) accurately and kept in a desiccators containing 50gm anhydrous calcium chloride. After three days, the formulations were weighed. The percentage moisture loss was calculated using the formula as follows: Percentage moisture loss = initial weight – final weight/final weight x100 the Percentage moisture loss = 1gm.

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366



International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



#### (ii) Transparency, smoothness and weight on drying:

The 5 ml gel formulation taken in the 10ml test tube and visually checked for its transparency. The smoothness of the gel formulation was tested by rubbing between the fingers and observes whether the gel is smooth, clumped, homogenous or rough. The relative density of the formulation or weight/ ml of the formulation was determined by taking the weight in gm. of 10 ml formulation and 10 ml distilled water using RD bottle.

### V) Viscosity pH and microbial growth:

IJARSCT

ISSN: 2581-9429

Viscosity is an important feature to determine the resistance of flow of gel formulation so that it can spread on the skin properly. It was determined with the help of viscometer (Brookfield) using 2 number spindles. pH of the formulation was determined by using pH meter (Elico). In this method, electrode was washed with double distilled water, dried with the help of tissue paper and then dipped in 20ml gel formulation. The average pH (n=3) of the gel formulations were recorded at ambient condition.

#### **II. CONCLUSION**

The present study is aimed to test the quality of Aloe Vera gel in multiple aspects. By using Aloe Vera gel, showed a multipurpose effect and all these ingredients showed significant different activities. Based on results and discussion, the formulations F1H, F2H and F3H were stable at room temperature and can be safely used on the skin.

Aloe Vera gel are used to stimulate blood circulation, rejuvenates the muscles and help to maintain the elasticity of the skin and remove dirt from skin pores. The advantage of gel is their nontoxic nature, reduce the allergic reactions and time tested usefulness of many ingredients.

The formulation was found homogenous, easily washable and also had very slightly alkaline pH which were compatible with normal skin physiology. Angle of repose is a characteristic related to inter particulate friction or resistance to the movement between the particles. The flow property has been classified as per limit of Indian Pharmacopoeia in terms of the angle of repose. The results of all these parameters indicated that the gel of combined form possess good flow properties and good packing ability. Consequently, it exhibited good flow properties for formulation to achieve soft, fresh and clean formulation.

### REFERENCES

1] Aggarwal D and Barna K. (2004). Tissue culture propagation of elite plant of Aloe Vera Linn. Journal of Biochemistry and Biotechnology, 13, 77-79.

2] Barna, K.S., Wakhlu ,A.K. 1994. Whole plant regeneration of Cicer arietium from callus culture via organogenesis. Plant Cell Reports. 13: 510-513.

3] S.AHMED,A.H. KABIR, M.B. AHMEDI,M.A. RAZVY2 and S. GANESAN [2007] An efficient method has been developed using shoot tip explants in Aloe vera.

4] N. Dwivedi, A. Indiradevi, K. Asha, N. Asokan, and A. Suma, "A protocol for micropropagation of Aloe Vera L. (Indian Aloe)-a miracle plant," Research in Biotechnology, vol.5, pp. 01-05, 2014.

5] Steenkamp V and Stewart MJ. (2007). Medicinal applications and toxicological activities of Aloe products. Pharmaceutical Biology, 45, 411-420.

6] Chand, S. and Roy, S.C., 1981. Induction of organogenesis in callus cultures of Nigella sativa L. Ann. Bot., 48: 1-4.

7] Durzan, D.J., 1984. Special problems: Adult vs. juvenile explants. In:W.R. Sharp,

D.A. Evand, P.V. Ammirato and Y. Yamada (Editors), Handbook of Plant Cell Culture. MacMillan, New York, pp. 47-63.

8] Schaik AH, Van Struik PC, Damian TG. Effects of irrigation and N on the vegetative growth of Aloe barbadensis Mill-in Aruba. Trap Agric 1997;74(2):104-9. 27.

9] Natali L, Sanchez IC, Cavallini A. In vitro culture of Aloe barbadensis Miller micropropagation from vegetative meristems. Plant Cell Tiss. Org. 1990;20:71-74.

10] Meyer HJ, Staden JV Rapid in vitro propagation of Aloe barbadensis Miller. Plant Cell Tiss. Org.1991 ;26: 167-171.

Copyright to IJARSCT www.ijarsct.co.in



DOI: 10.48175/IJARSCT-27366





International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

# Volume 5, Issue 1, June 2025



11] Gage, D. (1996). Aloe vera: Natures Soothing Healer. Healing Acts Press, Rochester, Vermont, USA, 120 p. 3

12] Surjushe, A., Vasani, R., &Saple, D. G. (2008). Aloe vera: a short review. Indian journal of dermatology, 53(4), 163-166.

13] Schulz, V., Hansel, R., & Tyler, V. E. (1997). Rational Phyto therapy: A Physicians' Guide to Herbal Medicine. Berlin: Springer, 306.

14] Christaki, E. V., & Florou-Paneri, P. C. (2010). Aloe vera: a plant for many uses. J Food Agric Environ, 8(2), 245-249.

15] Yagi, A., Tsunoda, M., Egusa, T., Akasaki, K., & Tsuji, H. (1998). Immunochemical distinction of Aloe vera, A. Arborescens, and A. Chinensis gels. Plantamedica, 64(03), 277-278.

16] Van Wyk, B. E., Van Oudtshoorn, M. V. R., & Smith, G. F. (1995). Geographical variation in the major compounds Of Aloe ferox leaf exudate. PlantaMedica, 61(03), 250-253

17] Saito, M., Tanaka, M., Misawa, E., Yao, R., Nabeshima, K., Yamauchi, K., ...& Furukawa, F. (2016). Oral administration Of Aloe vera gel powder prevents UVB-induced decrease in skin elasticity via suppression of overexpression of MMPs in hairless mice. Bioscience, biotechnology, and biochemistry, 80(7), 1416-1424..

18] García, M. H., Juárez, J. A. T., & Jiménez, A. D. (2019). Importance and Properties of Aloe vera in the Production of Hair Shampoo. The Journal of Middle East and North Africa Sciences, 5(7), 18-23.

19] Guenther, L., Lynde, C. W., Andriessen, A., Barankin, B., Goldstein, E., Skotnicki, S. P.,...& Sloan, K. (2012). Pathway 2 / 4To dry skin prevention and treatment. Journal of cutaneous medicine and surgery, 16(1), 23-31

20] Komatsu, D., Mistura, D. V., Motta, A., Domingues, J. A., Hausen, M. A., &Duek, E. (2017). Development of a Membrane of poly (L-coD, L lactic acid-co-trimethylene carbonate) with aloe vera: An alternative biomaterial Designed to improve skin healing. Journal of biomaterials applications, 32(3), 311-320.

21] SharrifMoghaddasi, M., & Res, M. (2011). Aloe vera their chemicals composition and applications: A review. Int J Biol Med Res, 2(1), 466-471.

22] King, G. K., Yates, K. M., Greenlee, P. G., Pierce, K. R., Ford, C. R., McAnalley, B. H., & Tizard, I. R. (1995). The effect Of Acemannan Immunostimulant in combination with surgery and radiation therapy on spontaneous canine and Feline fibrosarcomas. Journal of the American Animal Hospital Association, 31(5), 439-447.

23], K., & He, Q. (2004). Aloe vera: a valuable ingredient for the food, pharmaceutical and cosmetic industries A review. Critical reviews in food science and nutrition, 44(2), 91-96. Dat, A. D., Poon, F., Pham, K. B., &Doust, J. (2012). Aloe vera for treating acute and chronic wounds. Cochrane Database of Systematic Reviews, (2).

24] Ghaffarzadegan, R., Alizadeh, S. A., Ghaffarzadegan, R., Haji Agaei, R., &Ahmadlou, M. (2013). Effect of aloe vera Gel, compared to 1% silver sulfadiazine cream on second- degree burn wound healing. Complementary Medicine Journal, 3(1), 418-428.

25] Sahu, P. K., Giri, D. D., Singh, R., Pandey, P., Gupta, S., Shrivastava, A. K., ...& Pandey, K. D. (2013). Therapeutic and Medicinal uses of Aloe vera: a review. Pharmacology & Pharmacy, 4(08), 599-610 3 / 4

26] Maenthaisong, R., Chaiyakunapruk, N., Niruntraporn, S., &Kongkaew, C. (2007). The efficacy of aloe vera used for Burn wound healing: a systematic review. Burns, 33(6), 713-718.

27] Surjushe, A., Vasani, R., &Saple, D. G. (2008). Aloe vera: a short review. Indian journal of dermatology, 53(4), 163166

28] Hamman, J. H. (2008). Composition and applications of Aloe vera leaf gel. Molecules, 13(8), 1599-1616.

29] Samarghandian S, Farkhondeh T, Samini F. Honey and Health: A Review of Recent Clinical Research. Pharmacognosy Res. 2017 Apr-Jun;9(2):121-127. Doi: 10.4103/0974- 8490.204647. PMID: 28539734; PMCID: PMC5424551.

30] Ranneh, Y., Akim, A.M., Hamid, H.A. et al. Honey and its nutritional and anti- inflammatory value. BMC Complement Med Ther 21, 30 (2021). https://doi.org/10.1186/s12906-020-03170-5

31] Safitri, Fenny & Nawangsari, Desy & Febrina, Dina. (2021). Overview: Application of Carbopol 940 in Gel. 10.2991/ahsr.k.210127.018.

32] USPNF Glycerin, The United States Pharmacopeial Convention, USP, Accessed Date: 01/09/2023. Available from: https://www.uspnf.com/sites/default/files/usp\_pdf/EN/USPNF/revisions/2009-01- 30glycerinmonograph.pdf

DOI: 10.48175/IJARSCT-27366









International Journal of Advanced Research in Science, Communication and Technology

International Open-Access, Double-Blind, Peer-Reviewed, Refereed, Multidisciplinary Online Journal

#### Volume 5, Issue 1, June 2025



33] Aslani A, Zolfaghari B, Fereidani Y. Design, formulation, and evaluation of a herbal gel contains melissa, sumac, Licorice, rosemary, and geranium for treatment of recurrent labial herpes infections. Dent Res J (Isfahan). 2018
May-Jun;15(3):191-200. Doi: 10.4103/1735-3327.231865. PMID: 29922338; PMCID: PMC5958536.4 / 4
34] Misal, G. & Dixit, G. & Gulkari, Vijay. (2012). Formulation and evaluation of herbal gel. Indian Journal of Natural Products and Resources. 3. 501-505.



DOI: 10.48175/IJARSCT-27366

