

# A Review -Synergistic Anticancer Potential of Ginger, Bloodroot, and Aloevera: Development and Mechanistic Evaluation of A Polyherbal Formulation

**Ms. Pratibha J. Kale<sup>1</sup>, Dr. Kazi Mehraj Abukalam<sup>2</sup>, Dr. Annasaheb B. Jagnar<sup>3</sup>**

Shri JTT University, Jhunjhunu, Rajasthan, India<sup>1,2</sup>

Amrutvahini Institute of Pharmacy, Sangamner, MS, India<sup>3</sup>

**Abstract:** *The present study focuses on the optimization and standardization of a polyherbal anticancer tablet formulation using three medicinal plants: Aloe vera, Bloodroot, and Ginger, selected for their known anticancer properties. Phytochemical analysis revealed the presence of flavonoids, alkaloids, phytosterols, and phenolic compounds in the plant extracts. The physical properties of the tablet granules, including angle of repose, bulk density, Hausner ratio, and compressibility index, indicated good flow characteristics. Tablets demonstrated uniformity in weight, adequate hardness, mechanical stability, and appropriate disintegration times within pharmacopoeial limits. Stability studies over three months showed no significant deviations in key parameters, confirming the formulation's stability. Acute toxicity tests, conducted in line with WHO and OECD guidelines, indicated high LD50 values, affirming the safety of the plant extracts. The in vivo anticancer activity was tested at concentrations of 200 and 400 mg/kg, with histopathological studies revealing the effect on solid tumors in various tissues. The polyherbal tablet formulation, developed on a laboratory scale, shows promise as a stable, cost-effective dosage form for potential use in cancer management, combining traditional medicinal knowledge with modern pharmaceutical techniques*

**Keywords:** Bloodroot, Aloe vera, Ginger, Polyherbal

## I. INTRODUCTION

Plants have a unique position in the universe since they are the foundation of all life on Earth. They are the main producers in every food chain. Humans get 80% of their protein and 90% of their calories directly from plants. People have used plants as potential sources of medicine since the dawn of humanity. India is the source of the abundance of medicinal herbs, however traditional medicine is still practiced there. Therefore, it is imperative that this enormous natural resource be improved and used in accordance with the needs of humans and the growth of technology.

Cancer is one of the primary human diseases that causes huge suffering and economic loss world-wide. Chemotherapy is among the cancer treatments available. Unfortunately, the side effects of chemotherapy medications can be fatal and they are poisonous. New methods in the fight against cancer and its treatment in humans are constantly in the works. Natural ingredients provide the basis for around 60% of the anticancer drugs currently available. A large number of currently used medications have natural products as their primary ingredient. Certain plants and herbal items include phytochemicals that may be used as therapeutic or preventative agents against different types of cancer in humans. The cell either gets rid of or stores the byproducts of the metabolic process when it absorbs chemical carcinogens. The cellular metabolism of carcinogens and their byproducts can influence the expression and functioning of genes in a variety of ways, including cell cycle control, DNA repair, development of cells, and cell death. A number of genotoxic processes, including chromosomal cracking, fusion, deletion, mis-segregation, and non-disjunction, are induced by some carcinogens. Iron, arsenic, and cadmium ions or molecules can cause chromosomal variations in quantity and



structure. Some of these agents suppress the immune system, while others trigger inflammation, activate receptors like arylhydrocarbon or oestrogen receptors, generate reactive oxygen species, or even silence epigenetic factors. Hypermutability, a genome-wide loss of control over proliferation, immunity to apoptosis, and other hallmark traits of cancer cells may occur as a result of alterations to signal-transduction processes brought about by a mix of genotoxic and nongenotoxic causes.

## **II. PLANT PROFILE**

### **1. ALOEVERA**

**Synonym:** Aloe vulgairis Lamarek, or Aloe barbadensis Mil. or Aloe officinalis Forskal.

**Botanical Classification:**

**Kingdom:** Plantae

**Division:** Angiosperms

**Class:** Monocots

**Order:** Asparagales

**Family:** Asphodelaceae

**Genus:** Aloe

**Species:** Aloe vera

### **Biological Source**

Dried juice extracted from the undersides of the leaves of several varieties of Aloe plants is known as aloe vera. Members of the family Liliaceae, including Aloe perryi Miller, Aloe vera Linn, Aloe barbadensis Mil, and Aloe ferox Miller.

### **Geographical Source**

Native to the Arabian Peninsula, Aloe vera is now widely cultivated in tropical, subtropical, and arid regions across the world. Countries like India, China, Mexico, the United States, and parts of Africa are notable for Aloe vera cultivation.

### **Plant Description**

**Type:** Perennial, succulent.

**Size:** Typically grows 60–100 cm (24–39 inches) in height.

#### **Leaves:**

Fleshy, thick, lance-shaped.

Greenish or gray-green with serrated edges.

Can grow up to 60 cm long and 8 cm wide at the base.

#### **Flowers:**

Yellow or orange, tubular in shape, occurring in a spike up to 90 cm tall.

Blooms during the summer.

**Roots:** Shallow, fibrous root system.


**Stem:** Short, sometimes nearly stemless.

### **Chemical Constituents**

The three key components of aloes, namely  $\beta$ -barbaloins, are aloin isomers, Isobarbaloins, and Barbaloin. These three isomers form what is known as "crystalline" aloin and are to varying degrees (10% to 30%) throughout the course of treatment. Additionally, there are components such as resin, emodin, Aloe-emodin, and amor-phous aloin. All of the cultivars include barbaloin, a crystalline glycoside that has a somewhat yellow color, is bitter, and is soluble in water. Crystalline isobarbaloins are found in aloe from Curacao, in trace amounts in aloe from Cape Cod, and missing from aloe from Socotrine and Zanzibar.  $\beta$ - and Barbaloin are the main ingredients of Zanzibar aloe and Socotrine.



## 2. GINGER



### Synonyms

Rhizoma zingiberis, Zingibere.

### Biological Source

The dehydrated rhizomes of the *Zingiber officinale* Roscoe plant, which is a member of the Zingiberaceae family, are what make ginger.

### Origin and distribution

Africa, Nigeria, Jamaica, India, Japan, and the West Indies are the primary places where it is grown. Southeast Asian maritime regions are the source of ginger. It does not exist in the wild; it is a real cultigen. The earliest known evidence of its domestication dates back to the Austronesian peoples, who have long farmed and used several types of ginger. Other varieties of ginger were grown by them, such as bitter ginger (*Zingiber zerumbet*), white turmeric (*Curcuma zedoaria*), and turmeric (*Curcuma longa*). The leaves and rhizomes were either eaten raw or used to flavor dishes. Mats were also woven from the leaves. Aside from these applications, Austronesians considered ginger to be sacred, using it in healing rituals and in prayers to spirits for protection.

### Characteristics

The rhizomes measure roughly 1.5 cm in thickness, 3 to 6 cm in width, and 5 to 15 cm in length. Branches of the Jamaica ginger are found. Its outer surface is buff yellow in color with longitudinally striated fibers, and it displays sympodial branching. The section of the buds has small circular depressions, and the broken surface displays a well-developed endodermis, a wide stele, and a slender bark with sporadic little points of fibrovascular bundles and yellowish secretion cell points. The ginger tastes strong and pleasant and has a fragrant, pleasant smell.

### Chemical Constituents



The distinct aroma and taste of ginger stem from its volatile oils, which make up 1-3 percent of the weight of fresh ginger. These oils are mainly composed of sesquiterpenes, which include zingerone, beta-bisabolene, zingiberene, shogaols, and gingerols. The main aromatic compound in ginger is -gingerol (1- [4'-hydroxy-3'- methoxyphenyl]-5-hydroxy-3-decanone). The chemical composition of raw ginger is about 400.

During the drying process, gingerols yield zingerone, which has a spicy-sweet scent and less pungency. Shogaols, which are more aromatic, are made from gingerols by heating, storing, or by acidity. Other components include a variety of compounds with monoterpenes, as well as amino acids, protein, fibre, phytosterols, vitamins, and minerals. Additionally, fresh ginger contains zingibain, an enzyme that catalyses cysteine proteases similar to rennet.

Ginger comprises starch, 5–8% pungent resinous substance, and 1–2% volatile oil. The medicine's pleasant perfume comes from the volatile oil, and its strong flavour comes from the odourless, yellowish-oily gingerol. Sesquiterpene hydrocarbons such as  $\alpha$ -zingiberol,  $\alpha$ -bisabolene,  $\alpha$ -farnesene, and  $\alpha$ -sesquiphellandrene, as well as  $\alpha$ -sesquiterpene alcohol, make up volatile oil.

Some ingredients, such as shogaol and zingerone, are less strong. Fresh rhizomes do not contain shogal, which is a byproduct of gingerol's dehydration.





### 3. BLOODROOT

**Synonym:** Bloodroot; Bloodworth, Red Root, Red Puccoon

#### Biological Source

The poppy family Papaveraceae counts this plant as the sole member of the genus *Sanguinaria*.

#### Geographical Source

This wildflowers, which blooms in the spring, is native to North America and can be originate in verdant meadows from Nova Scotia and Florida westward to Nebraska, Manitoba, and Alabama. The permanent herbaceous flowering plant called bloodroot originates in the eastern part of North America. From the Atlantic coast to the Gulf of Mexico and beyond, westward to the Great Lakes and the Mississippi River embayment, bloodroot can be located throughout eastern North America.

#### Cultivation

*Sanguinaria canadensis* is grown for its aesthetic qualities. Gardeners appreciate the huge, brilliant white flowers that the double-flowered varieties produce quite early in the gardening season. Bloodroot flowers have a brief blooming show since their petals shed within a day or two of fertilization, while their double versions bloom for much longer. Because the stamens in the double blooms have been transformed into components resembling petals, pollination is more challenging.

#### Characteristics

The height of bloodroot ranges from 20 to 50 cm (8 to 20 in). It has one enormous basal leaf with five to seven lobes that can reach a diameter of up to 25 cm (10 in). Bright orange to crimson sap shoots from a reddish rhizome from which the leaves and flowers emerge. The genus name *Sanguinaria* (from the Latin *sanguinarius*, "bloody") derives from the color of the sap. Every year, the rhizomes lengthen and divide into colonies. In early spring, plants begin to bloom before the foliage appears. The leaves unfold to their full size following blossoming. Mid- to late-summer is when plants go dormant, It happens later than other fleeting spring flowers.

#### Chemical Constituents

When cut, bloodroot releases a brilliant orange-red sap from all parts of the plant; the rhizome, which stores the juices, has the largest concentrations. The sap is rich in starch, scarlet resin, and isoquinoline alkaloids. The phytochemicals. Isoquinoline alkaloids, primarily sanguinarine and chelerythrine, are abundant in sanguinaria root. Unlike most other alkaloids, sanguinarine, a benzophenanthridine alkaloid, exhibits a red color in aqueous solutions.

Isoquinoline alkaloids, primarily sanguinarine and chelerythrine, are abundant in sanguinaria root. Unlike most other alkaloids, sanguinarine, a benzophenanthridine alkaloid, exhibits a red color in aqueous solutions. The rhizomes and roots contain the highest and second-highest concentrations of it, respectively, with lower concentrations found in leaves and flowers. Protopine and berberine are two related chemicals found in the plant, along with other minor alkaloids.

### III. PREPARATION OF CRUDE DRUG FOR EXTRACTION

The extract was made from certain plant leaves. Gathered and left to dry in the shade were specific plant leaves. The leaves were dried and then finely ground in a machine grinder. Then, sieve No. 16 was used to screen this coarse



powder made from the leaves of various chosen plants. Additionally, they were kept for the extraction in an airtight container after passage.

To extract the desired elements from the unrefined drug, Soxhlet extraction is utilized.

#### **Aloevera**

Water was used as the solvent throughout the 48-hour extraction of 50 grams of dried coarse powder from Aloevera utilizing a Soxhlet equipment. After complete extraction, the aqueous extract was concentrated in a vacuum dryer at 40°C under reduced pressure to preserve it in desiccators.

#### **Bloodroot**

50 grams of dried coarse Bloodroot powder were continuously extracted with water as the solvent for 48 hours using a Soxhlet device. After complete extraction, the aqueous extract was concentrated in a vacuum dryer at 40°C under reduced pressure to preserve it in desiccators.

#### **Ginger**

Water was used as the solvent in a Soxhlet device to continuously extract 50 grams of dried coarse powder gingerwa over the course of 48 hours. Desiccators were used to store the dried extract that was created by concentrating the solution of water under less pressure at 40°C in a vacuum drier after extracting.

#### **Extraction**

Extraction of three plants was prepared individually as well as in combinations of 2:1:1 and 4:1:1 by continuous hot extraction using Soxhlet extraction method for 48 hrs. Physical examination was performed on the extracts (color, consistency).



**Figure No. 5.1: Soxhlet Extraction of Aloevera**



**Figure No. 5.2: Soxhlet Extraction of Ginger**

#### **IV. CONCLUSION**

The Global Effects of Cancer and Complementary Medicine's Role

Copyright to IJAR SCT  
[www.ijarsct.co.in](http://www.ijarsct.co.in)



DOI: 10.48175/568





With its increasing incidence and fatality rates, cancer continues to rank among the world's most pressing health concerns. This has led to a rise in interest in complementary medicine, which emphasizes the use of herbal remedies in addition to traditional therapy. The possible anticancer benefits of polyherbal preparations, which combine many medicinal herbs, have drawn attention among these alternative medicines. Because these formulations have therapeutic uses in conventional medical systems, they are widely employed.

Developing a polyherbal tablet product is the primary objective of this research with an anticancer focus. In order to guarantee effectiveness, safety, and stability, the main objectives are to improve the formula and standardize its preparation. Research in this study looks into the potential anti-cancer properties of the selected plants and the stability of the formulation as it is turned into a solid dosage form (tablet).

The following three medicinal plants were selected for assessment of their anticancer potential after a thorough analysis of the literature:

- Aloe vera (Aloe barbadensis miller)
- Bloodroot (Sanguinaria canadensis)
- Ginger (Zingiber officinale)

These plants were chosen because they are already in use in medical settings and because they may have anticancer properties. Bioactive substances found in all plants, including phytosterols, alkaloids, flavonoids, and phenolic compounds, possess numerous demonstrated therapeutic applications, such as the ability to fight cancer and act as antioxidants.

**Aloe vera:** Aloe vera includes chemicals that are known to have therapeutic qualities like aloesin and aloesin derivatives, which have shown anticancer and antioxidant effects.

**Bloodroot:** This plant contains alkaloids, particularly sanguinarine, which looked into because of its capacity to cause cancer cells to undergo apoptosis, often known as the programmed death of cells.

**Ginger:** Ginger contains chemicals such as gingerol and shogaol, which have anti-inflammatory and antioxidant activities, which are thought to help in inhibiting cancer cell proliferation and promoting cell death in tumors.

## REFERENCES

- [1]. Abiy E and Berhe A. Anti-bacterial effect of Garlic (*Allium sativum*) against clinical isolates of *Staphylococcus aureus* and *Escherichia coli* from patients attending Hawassa Referral Hospital, Ethiopia. *J Infec Dis Treat.* 2016; 2(2): 18.
- [2]. Abutaha NM, Farooq MF, Mohammed AZ, Alotaibi A, Cordero MA, Bepari A, Alarifi S. Cytotoxic activity and toxicity study of HF8, a poly-herbal formulation. *Journal of King Saud University-Science.* 2021;33(3):101377.
- [3]. Adebayo I A, Balogun W G and Arsad H. *Moringa oleifera*: An apoptosis inducer in cancer cells. *Trop J Pharma Res.* 2017;16 (8): 2289-2296.
- [4]. Ahmad R, Ahmad N, Naqvi A A, Shehzad and Ghamdi M S. Role of traditional Islamic and Arabic plants in cancer therapy. *J Tradi Complem Med.* 2017;7(2): 195- 204.
- [5]. Ahmedin Jemal, Rebecca Siegel, Elizabeth Ward, Taylor Murray, Jiaquan Xu, Michael J. Thun. Cancer Statistics. *Cancer J Clin.* 2007, 57: 43–66.
- [6]. Ahmedin Jemal, Rebecca Siegel, Elizabeth Ward, Taylor Murray, Jiaquan Xu, Michael J. Thun. Cancer Statistics. *Cancer J Clin.* 2007, 57: 43–66.
- [7]. Akev N, Candoken E, Erdem Kuruca S. Comparative Study on the Anticancer Drug Potential of a Lectin Purified from Aloe Vera and Aloe- Emodin. *Asian Pac J Cancer Prev.* 2020;21(1):99-106.
- [8]. Akimoto M, Iizuka M, Kanematsu R, Yoshida M, Takenaga K. Anticancer Effect of Ginger Extract against Pancreatic Cancer Cells Mainly through Reactive Oxygen Species-Mediated Autotic Cell Death. *PLoS One.* 2015;10(5):e0126605.



- [9]. Almawlah Y H. Antibacterial Activity of Punica granatum, Allium sativum and Piper nigrum against methicillin resistant Staphylococcus aureus (MRSA) isolated from wound infections in Al-Hilla, Iraq. Inter J Chem Tech Res. 2017; 10(2): 604-611.
- [10]. Alwe JR, Alwe RS. Pharmacognostic study of Strychnos potatorum Linn- A review. Int Ayurvedic Med J 2016;4(7):1222-6.
- [11]. Annappan U, Kumudhavalli MV, Mohan K. Development and Statistical Optimization of Polyherbal Tablets Containing Indigenous Plant Extracts. International Journal of Advancement in Life Sciences Research. 2024; 7(3):98-108.
- [12]. Annappan, Umesh. Formulation And Evaluation of Polyherbal Formulation Containing Indigenous Medicinal Plants. Section A-Research paper Eur.2023. 3719-3726.
- [13]. Annappan, Umesh. Formulation And Evaluation of Polyherbal Formulation Containing Indigenous Medicinal Plants. Research paper Eur. 2023; 3719- 3726.
- [14]. Aquil F, Ahmad I, Mehmood Z, “Antioxidant and free radical scavenging properties of twelve traditionally used Indian medicinal plants”, Turkish Journal of Biology, 2006; 30: 177-183.
- [15]. Arora D S and Kaur G J. Antibacterial activity of some Indian medicinal plants. J Natur Med. 2007; 61(3): 313–317.

