

Development and Evaluation of Herbal Wound Healing Formulations

Mr . Pramod B. Chikkodi, Tanuja Katkar, Rutuja A. Pujari, Sanika T. Hubale, Mahima P Magdum
Nootan College of Pharmacy, Kavathe Mhankal.

Abstract: The aerial parts of *Centella asiatica* (L.) Urban and rhizomes *Curcuma longa* (L.) were collected, dried under shade and extracted with ethanol and water respectively. These extracts were dried. Aloe barbadensis concentrated gel powder was obtained as free sample. Each extract and Aloe gel was added with appropriate proportion in cream base and evaluated for wound healing activity against povidone iodine ointment. In excision wound model the complete wound healing was observed with cream formulation I and II treated rats was observed in 18 days where as povidone iodine ointment took 16 days. In incision wound model tensile strength of formulations I and II was found to be 366.08 ± 2.32 and 351.35 ± 3.29 gm respectively. Tensile strength of povidone iodine ointment was found 379.98 ± 2.95 gm. Herbal formulation I and II took two days more for complete wound healing as compared to povidone iodine ointment.

Keywords: Centella asiatica, Cream formulation, wound healing activity

I. INTRODUCTION

Wound is the disruption of cellular and anatomic and functional continuity of living tissue, produced by physical, chemical, electrical or microbial insults to the tissue¹. Wound healing is the dynamic process take place by regeneration or repair of broken tissue².

Normal wound-healing response begins with injury and is a concerted sequence of events. The healing cascade is activated when platelets come into contact with exposed collagen leading to platelet aggregation and the release of clotting factors resulting in the deposition of a fibrin clot at the site of injury. The fibrin clot serves as a provisional matrix and sets the stage for the subsequent events of healing³. Inflammatory cells also arrive along with the platelets at the site of injury providing key signals known as cytokines or growth factors⁴. The fibroblast is the connective tissue responsible for collagen deposition that is needed to repair the tissue injury⁵. In normal tissues, collagen provides strength, integrity, and structure. When tissues are disrupted following injury, collagen is needed to repair the defect and restore anatomic structure and function⁶. In the present work formulations were prepared to accelerate wound healing process.

Literature survey revealed that *Centella asiatica* has the antimicrobial⁷, antioxidant activity⁸, wound healing property⁹ and increase collagen synthesis and tensile strength¹⁰. *Curcuma longa* has anti-inflammatory, antimicrobial¹¹ and wound healing property¹². Aloe has the anti-inflammatory activity¹³, increase degree of cross-linking¹⁴. Aloe also provides moisturizing activity.

In present study for easy application on wound o/w cream were prepared and their feasibility was checked by wound healing activity in wistar rats as compared to povidone iodine ointment.

II. MATERIAL AND METHOD

Preparation of extract

Centella asiatica (L.) Urban and *Curcuma longa* (L.) were collected from Amravati. The Plant species were identified and confirmed by Botanist Dr.Prabha Bhogaonkar, Director, Govt. Vidarbha institute of science and humanities, Amravati. A herbarium was prepared and deposited at Dept. of Botany, Govt. Vidarbha institute of science and humanities, Amravati.



Air-dried aerial parts of *Centella asiatica* (L.) Urban were defatted with petroleum ether and extracted with 90% ethanol in Soxhlet apparatus for 8 hours. The extract was concentrated on water bath and dried at 50⁰ □ 5⁰C in hot air oven. The powder of *Curcuma longa* rhizomes was boiled in purified water for 2 hours. The solution was then filtered through a coarse sieve twice.

Finally, the filtrate was concentrated on water bath to a thick paste like consistency and dried at 50⁰ □ 5⁰C in hot air oven. *Aloe barbadensis* concentrate gel powder was obtained as a free sample from Chaitannya Biologicals Private Limited, Malkapur (M.S.).

Preparation of Formulation

Table No. 1 Formulation I and II

Ingredients	Formulation I quantity (g)	Formulation II quantity (g)
<i>Centella asiatica</i> extract	6	5
<i>Curcuma longa</i> extract	1	0.5
<i>Aloe barbadensis</i> Gel	2	2
Polysorbate 60	5	5
White Soft Paraffin	25	25
Cetosteryl alcohol	4	4
Glycerin	12	12
Methyl paraben	0.1	0.1
Propyl paraben	0.05	0.05
Butylated hydroxyl anisole	0.02	0.02
Purified water	q.s.to 100 gm	q.s.to 100 gm

Procedure

Aloe barbadensis concentrate gel powder was added in purified water and kept overnight. Phase I was prepared by dissolving *Centella asiatica* extract, *Curcuma longa* extract and above prepared aloe gel in purified water and previously heat to then glycerin was mixed the solution with constant stirring.

Phase II was prepared by melting white soft paraffin at 70⁰C, then cetosteryl alcohol and polysorbate 60 was added to it with constant stirring. Lastly Methyl and propyl paraben and Butylated hydroxyl anisole (BHA) were added the mixture was stirred.

Phase II was added to Phase I with constant stirring at 70⁰C. Then allow cooling with constant stirring.

II. EVALUATION OF WOUND HEALING ACTIVITY

The experimental protocol was approved by institutional animal ethical committee of Anuradha College of Pharmacy, Chikhli Dist. Buldana as per guidelines issued by Committee of Purpose of Control and Supervision of Experiment on Animals (CPCSEA) India.

Animals

Male wistar rats (150-200g) were used for study. All animals had free access to pelleted food and water *ad libitum*. Temperature was maintained at 23± 1⁰C.

Treatment

Animals were wounded under light ether anesthesia, semiseptically. The animals were assigned into five group (n=6). Group I was untreated group, this was taken as control. Group II animals were received Povidone iodine ointment



treatment. Group III animals was received Cream base treatment, while groups of IV and V animals were received test formulations I and II treatments respectively in both excision and incision wound models. No other topical or systemic therapy was given to animals during the course of this study.

Excision wound model¹⁵

Hairs were removed from dorsal thoracic central region of anaesthetized rats. Full thickness from the demarked area was excised to produce wound measuring around 300 mm². Wound was cleaned with cotton swab soaked in alcohol. The two test formulations, cream base and povidone iodine ointment were applied on wound once daily for 20 days starting from the first day of wounding. Wound contraction was measured for 20 days at interval of 2 days.

Incision wound model¹⁶

Animals were anaesthetized and paravertebral incisions (2.5-3.0 cm long) were made through the entire length of skin. After the incision was made, the parted skin was kept together and stitched with nylon thread at 0.5 cm apart with curved needle (No. 11). The two test formulations, cream base and povidone iodine ointment were applied on wound once daily for 7 day. The sutures were removed on day 8 and wound tensile strength was measured on day 10 by using constant water flow technique.

Constant water flow technique

On the 10th day the animals was secured to the operation table, under light ether anaesthesia. A line was drawn on normal skin on either side of wound, 3 mm away from the wound line. Two Allis forceps were firmly applied on the lines facing each other. On one side the forceps was hooked firmly to metal rod fixed to the operation table. The other forceps was connected to a leakproof graduated polythene container through a string running over a pulley. The polythene container was connected to water reservoir placed at suitable height through a rubber tube kept occluded with a pinchcock. To measure wound tensile strength, the tube was released to allow a constant and continuous flow of water from the reservoir in to the polythene container. As the weight gradually increases, it acts as a pulling force to disrupt the wound. As soon as the gapping of the wound was observed, the rubber tube was clamped and the polythene container was weighed

Table 2 Excision wound model

Post woun -ding days	Wound Area(mm ²)				
	Control (Group I)	Standard (Group I)	Cream base (Group I)	Formulation I (Group I)	Formulation II (Group I)
0	330.58 ±18.20 (0)	325.11 ±10.51 (0)	334.86 ±18.13 (0)	324.06 ±13.25 (0)	327.80 ±17.66 (0)
2	270.83 ±4.30 (18.07)	229.58 ±3.35 (29.38)	266.98 ±2.22 (20.27)	234.65 ±6.20 (27.59)	239.90 ±5.46 (26.81)
4	248.03 ±4.72 (25.03)	195.88 ±4.83 (39.74)	236.31 ±5.23 (29.43)	271.26 ±2.13 (32.95)	221.90 ±2.01 (32.30)
6	232.03 ±1.72 (30.17)	178.38 ±1.89 (45.13)	225.48 ±5.06 (32.66)	182.65 ±2.25 (43.63)	183.28 ±2.45 (44.39)
8	194.71 ±2.59 (41.10)	129.73 ±2.32 (60.09)	187.61 ±1.50 (43.97)	135.21 ±2.94 (58.27)	145.85 ±2.41 (55.50)
10	184.60 ±2.53 (44.15)	100.58 ±2.36 (69.06)	174.95 ±3.20 (47.75)	114.85 ±2.07 (64.55)	122.00 ±1.63 (62.78)
12	169.25	63.88	149.83	79.58	87.53



	± 2.15 (48.80)	± 1.64 (80.35)	± 1.56 (55.25)	± 1.65 (75.44)	± 3.41 (73.29)
14	152.96 ± 2.12 (53.72)	21.5 ± 1.59 (93.38)	124.01 ± 3.17 (62.96)	36.73 ± 2.60 (88.66)	58.11 ± 1.75 (82.27)
16	122.30 ± 3.44 (63.00)	0(100)	107.10 ± 2.45 (68.01)	17.08 ± 2.36 (94.72)	26.45 ± 1.66 (91.93)
18	108.65 ± 2.70 (67.13)	0(100)	74.06 ± 2.12 (77.88)	0(100)	0(100)
20	93.53 ± 1.57 (71.68)	0(100)	56.46 ± 1.79 (83.13)	0(100)	0(100)

Value are mean $\pm$ SEM of Six animals (n=6) in each group. Number in parenthesis indicates percentage of wound contraction. All are significant at $p < 0.05$ as compared to group I (control) and indicate not significant.

Table No. 3 (Incision wound model)

Group	Treatment	Tensile strength(gm)
1	Control (Group I)	164.55 $\pm$ 3.14
2	Povidone iodine ointment (Group II)	379.98 $\pm$ 2.95
3	Cream Base (Group III)	202.75 $\pm$ 3.04
4	Formulation I (Group IV)	366.08 $\pm$ 2.32
5	Formulation II (Group V)	351.35 $\pm$ 3.29

Values are mean $\pm$ SEM of six animals in each group $p < 0.05$ Vs group I

III. RESULTS

In excision wound study (see table no.2), wound contraction progresses identically with Povidone iodine ointment and in wound treated with formulation I and II. In these three groups complete healing was observed between 16th and 18th day. While untreated group I (control) and group III (cream base) animals took more than 20 days for healing of wounds.

In incision wound study (see table no. 3), tensile strength of Povidone iodine ointment was found 379.98 $\pm$ 2.95 gm and group I (control) was found 164.55 $\pm$ 3.14 gm, while the group III (cream base) showed 202.75 $\pm$ 3.04 gm. Tensile strength of formulations I and II showed 366.08 $\pm$ 2.32 and 351.35 $\pm$ 3.29 gm respectively.

Statistical analysis

The results were expressed as mean $\pm$ SEM. The significance of differences between the means was analyzed by student's t- test followed by Turkey's test. A P-value <0.05 was considered significant.

IV. DISCUSSION

In the present work, plants like *C. asiatica*, *C. longa* and *A. barbadensis* were selected which accelerate wound healing process. The formulation I and II showed best results this may be due to asiaticoside from *C. asiatica* accelerate collagenisation and tensile strength¹⁰.

Asiaticoside, a major constituent of the herb is know to promoted wound-healing by reducing lipid peroxide levels in wounds⁹. Curcumin from *C. longa* is responsible for closure of wound and increase in collagen synthesis¹².

A. barbadensis increased degree of cross-linking as seen by increased aldehyde content and decreased acid solubility¹⁴. These all factors may be responsible for wound healing activity of formulations I and II.



In incision wound study, formulations enhanced the tensile strength of 10 days old wounds as compared with wound of untreated group. The importance of cross linking between collagen molecules and physical weave of collagen fibers in contributing to the tensile strength of wound is well acknowledged. Increase in tensile strength may be due to increase in collagen concentration per unit area and stabilization of fibers. Formulations I and II showed significant increase in tensile strength of the 10 day old wound. This may be due to promotion of collagenation and cross linking.

REFERENCES

- [1]. Schilling J.A. wound healing. *Psychological review*; 48(2): 374-415.
- [2]. Flanagan M. *Wound management* 1st edition. New York, Churchill Livingstone Inc., 22-26.
- [3]. Clark RA. Fibrin and wound healing. *Ann N Y Acad Sci*, 2001; 936: 355-67.
- [4]. Lawrence WT, Diegelmann RF. Growth factors in wound healing. *Clin Dermatol*, 1994; 12(1); 157-69.
- [5]. Ross R. Wound healing. *Sci Am* 1969; 220(6): 40-50.
- [6]. Prockop DJ, Kivirikko KI. Collagens: molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem* 1995; 64: 403-34.
- [7]. Oyediji OA and Afolayan AJ. Chemical Composition and Antibacterial Activity of the Essential Oil of *Centella asiatica* growing in South Africa. *Pharmaceutical Biology*, 2005, 43 (3) : 249-252
- [8]. Kumar MHV and Gupta YK. Effect of *Centella asiatica* on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clinical and Experimental Pharmacology and Physiology*, 2003, 30: 336-342
- [9]. Shukla A, et al. *J Ethnopharmacol*, 1999, 65(1): 1-11.
- [10]. Suquna L., Sivakumar P, and Chandrakasan G. Effect of *Centella asiatica* extract on dermal wound healing in rats, *Indian J Exp Biol.*, 1996 Dec, 34(12): 1208-11
- [11]. Ukil A, Maity S, Karmakar S, Datta N, Vedasiromoni JR. and Das PK. Curcumin, the major component of food flavour turmeric, reduces mucosal injury in trinitrobenzene sulphonic acid-induced colitis. *British Journal of Pharmacology*. 2003, 139: 209-218
- [12]. Das, suman k. (Jackson, MS), Use of turmeric in wound healing- United States patent 5401504.
- [13]. Kostalova D, et al. *Ceska Slov Farm*, 2004, 53(5): 248.
- [14]. Chithra P, Sajithal GB and Chandrakasan G., Influence of aloe vera on collagen characteristics in healing dermal wound in rats, *Mol Cell Biochem.*, 1998 Apr, 181(1-2): 71-76.
- [15]. International Journal of Advanced Research in Science, Communication and Technology : A review on Formulation and Evaluation of Natural Lip Balm by using Rose Petals
- [16]. Journal of Advancement in Pharmacognosy, 2025 Advanced Herbal Technology: A Review of Plant Identification, Authentication, and Advanced Extraction Techniques
- [17]. International Journals of Advanced Research in Science: A review on Formulation and Evaluation of Antifungal cream.
- [18]. Morton JJ and Malone MH, Evaluation of vulnerary activity by an open wound procedure in rats. *Arch Int Pharmacodyn*, 1972; 196, 117-26.
- [19]. Potawale SE, Yeole DU, Deshmukh Rani, Patil VG, Khopade AN and Purohit AP. Design, Development and evaluation of herbal wound healing formulation, *Adv. Pharmacol. Toxicol*, Vol 8 (3) 2007, 99-103

