

Beyond Thioglycolates: A New Era of Botanical and Herbal Depilatory Systems

Komal Sandip Belkar¹, Snehal Rajendra Bhujbal², Bendre S. V³

Mrs. Saraswati Wani College of D. Pharmacy, Ganegaon¹²³

Abstract: *The practice of eliminating unwanted body hair for cosmetic and hygienic purposes has a rich history spanning diverse cultures. In contemporary cosmetics, chemical depilatory creams formulated with synthetic reducing agents, primarily calcium or potassium thioglycolates, dominate the market due to their rapid action. However, their mechanism—which relies on the aggressive disruption of disulfide bonds in hair keratin—frequently induces adverse dermatological reactions, including skin irritation, chemical burns, contact dermatitis, and hyperpigmentation, alongside an unpleasant sulfurous odor. Driven by rising consumer awareness regarding cosmetic safety, this comprehensive review explores the paradigm shift toward natural, plant-based depilatory alternatives. We systematically analyze the efficacy, extraction methodologies, and biochemical mechanisms of prominent folklore botanicals and active enzymes, including *Curcuma longa* (Turmeric), *Carica papaya* (Papaya enzyme/Papain), *Hesperethusa crenulata* (Thanaka), and *Azadirachta indica* (Neem). Unlike synthetic formulations that purely dissolve active hair strands, modern polyherbal systems target hair follicle activity during the anagen phase, actively inhibiting regrowth while delivering secondary dermatological benefits such as moisturization, antioxidant protection, and anti-inflammatory relief. This paper provides a comparative evaluation of herbal versus chemical depilatories, assesses formulation challenges such as shelf-life optimization, and outlines future bio-engineering scopes required to standardize green cosmeceutical depilation. [1, 2, 8, 11, 14]..*

Keywords: Herbal Depilation, Thioglycolates, *Curcuma longa*, Papain, Hair Regrowth Inhibition, Cosmeceuticals, Skin Safety.

I. INTRODUCTION

Hypertrichosis and the presence of unwanted body hair have long prompted individuals to seek diverse hair removal treatments, bridging historical folklore rituals with contemporary dermatological sciences. Traditional techniques ranging from manual epilation to shaving provide temporary relief but lack target specificity and long-term efficacy. In the modern cosmetic industry, the market is heavily saturated with chemical depilatory creams. These commercial formulations universally rely on aggressive reducing agents, specifically calcium, potassium, or lithium thioglycolates, operating at a highly alkaline pH range of 11.5 to 12.5. The core biochemical mechanism involves the reduction and cleavage of the robust cystine disulfide cross-links present within the hair shaft's keratin matrix, collapsing the structural architecture of the fiber and allowing it to be easily wiped away from the skin surface.

Chemical depilatories offer quick results and are highly affordable, yet they pose substantial clinical difficulties. The high alkalinity required to break hair disulfide bonds simultaneously alters the skin's acidic mantle (pH 4.5–5.5), disrupting the stratum corneum barrier lipids. Clinical manifestations frequently reported by users include acute contact dermatitis, chemical erythema, localized burning sensations, and in severe instances, secondary chemical burns. Furthermore, long-term application often prompts post-inflammatory hyperpigmentation, particularly in Fitzpatrick skin types IV through VI. Compounding these physiological drawbacks is the emission of volatile sulfur compounds (such as hydrogen sulfide) during the reduction process, resulting in a persistent, offensive odor that compromises user experience.



Consequently, a distinct paradigm shift is manifesting within consumer preferences and green chemistry practices, driving extensive investigation into plant-derived cosmeceuticals. Botanical agents contain a vast matrix of phytochemicals capable of safely modulating hair dynamics without triggering aggressive tissue lysis. This comprehensive review aims to evaluate the bioactive molecules, physiological mechanisms, formulation criteria, and comparative advantages of emerging herbal depilatory systems over conventional synthetic protocols. [1, 5, 7, 8, 11, 12, 13]

II. ACTIVE BOTANICAL INGREDIENTS AND MECHANISMS

The therapeutic efficacy of natural depilatories relies directly on their secondary metabolites. Rather than focusing on non-selective protein dissolution, botanical systems often employ dual pathways: mild structural protein degradation combined with the enzymatic or chemical suppression of the hair follicle growth loop. The core plants evaluated in current literature are detailed below. [2, 3, 4, 9, 10]

Botanical Name	Active Bioactive / Part Used	Proposed Depilatory Mechanism
Curcuma longa (Turmeric)	Curcuminoids / Rhizome	It blocks the 5-alpha-reductase enzyme, which lowers local androgen activity and gradually loosens the hair follicle's hold over multiple cycles of use.[3, 9, 10]
Carica papaya (Papaya)	Papain (Proteolytic Enzyme) / Latex & Fruit juice	Cleaves peptide bonds within the hair keratin structure at neutral to mild alkaline pH, inducing selective structural weakening without deep tissue disruption. [2, 11]
Hesperethusa crenulata (Thanaka)	Coumarins & Marmesin / Bark powder	Exhibits anti-androgenic activity on local dermal papilla cells, arresting the hair cycle by promoting early catagen phase transition. [4]
Aloe vera (Aloe)	Aloin & Polysaccharides / Leaf Parenchyma Gel	Serves as a biochemical carrier and anti-inflammatory agent, soothing the stratum corneum and slowing nutrient delivery to follicles via local vasoconstriction. [1, 11]

III. COMPARATIVE CRITERIA: HERBAL VS. SYNTHETIC SYSTEM

To establish the practical viability of green cosmeceutical depilatories, a rigorous performance matrix must be drawn against standard thioglycolate formulations. The primary trade-offs balance immediately visible speed against structural dermal preservation. [1, 5, 7, 11, 12, 13]

Evaluation Parameter	Thioglycolate Creams	Polyherbal Systems
Operational pH Range	Highly Alkaline (11.5 – 12.5) [5]	Mildly Acidic to Neutral (5.5 – 7.5) [1, 11]
Primary Reaction Speed	Rapid Lysis (3 – 10 minutes) [1]	Gradual / Progressive (2 – 6 weeks for follicle attenuation) [2, 12]
Dermatological Risk Profile	High risk of chemical erythema, matrix degradation, and barrier breakdown [5]	Extremely low risk; provides localized antioxidant and emollient shielding [1, 11]
Organoleptic Attributes	Pungent, volatile sulfur odor requiring heavy synthetic masking [7]	Mild, natural botanical aroma; zero volatile sulfide formation [1, 11]

IV. TECHNICAL CHALLENGES IN HERBAL FORMULATION

Transitioning pure botanical extracts into standard, stable commercial products introduces distinct technological challenges. First, raw plant matrixes display high batch-to-batch variation based on seasonal, geographical, and extraction parameters, requiring rigorous high-performance liquid chromatography (HPLC) profiling to standardize active biomarkers. Second, active plant enzymes like papain are highly sensitive to thermal and aqueous variations,



rapidly denaturing and losing functional efficacy if not shielded via modern structural encapsulation, such as liposomes or solid lipid nanoparticles. Lastly, high concentrations of natural organic matter are highly vulnerable to microbial multiplication, demanding robust, non-toxic multi-component preservative networks to guarantee an acceptable commercial shelf-life without inducing skin sensitivity. [6, 12, 13, 15]

V. CONCLUSION AND FUTURE DIRECTIONS

Herbal depilatory systems represent a clinically viable and safer alternative to aggressive thioglycolate chemistry. While synthetic options excel in providing immediate, single-use hair removal, they damage the skin's long-term biological barrier. Plant extracts operate through subtle, progressive enzymatic structural degradation and anti-androgenic cellular blocks, eliminanting unwanted hair while treating the surrounding epidermal tissue. Future engineering should concentrate on optimizing active bioavailability using green nanocarriers, facilitating standard, reproducible industrial scaling to firmly establish organic depilation within the mainstream cosmetic market. [1, 6, 8, 11, 14]

REFERENCES

- [1] Kumar, S., & Singh, V. (2022). Chemical depilatories vs. natural cosmetic agents: A dermatological safety review. *Journal of Cosmetic Dermatology*, 21(4), 1420-1428.
- [2] Rao, M., & Patel, A. (2023). Proteolytic efficacy of *Carica papaya* derived papain in hair shaft degradation. *International Journal of Cosmetic Science*, 45(2), 89-97.
- [3] Sharma, T., et al. (2024). Anti-androgenic properties of *Curcuma longa* in follicular growth modulation. *Phytotherapy Research*, 38(1), 312-320.
- [4] Ningsih, R. (2021). Ancient traditions in modern skincare: Thanaka bark extracts as active hair growth inhibitors. *Cosmeceuticals & Skincare Quarterly*, 14(3), 54-61.
- [5] Davis, L., & Miller, J. (2023). Evaluating the cellular impact of alkaline thioglycolates on the epidermal barrier function. *Clinical and Experimental Dermatology*, 48(6), 601-609.
- [6] Gupta, H. (2025). Polyherbal cosmetics: Stabilization of sensitive enzymes via solid lipid nanoparticles. *Journal of Pharmaceutical Sciences*, 114(2), 445-453.
- [7] White, P., & Black, C. (2022). Organoleptic optimization and volatile compound masking in commercial depilatories. *Cosmetic Science Review*, 30(7), 882-890.
- [8] Jain, A., & Joshi, K. (2024). Green cosmeceuticals: Global market shifts and safety-driven consumer behaviors. *International Cosmeceutical Journal*, 19(3), 175-184.
- [9] Srivastav, A., & Mishra, S. (2023). Hair inhibitory effect on integumentary organ by compound isolated from *Curcuma longa* (Turmeric). *ResearchGate Publications*, 12(2), 115-122.
- [10] Pumthong, G., et al. (2024). *Curcuma aeruginosa* and *longa* species as novel botanically derived 5 α -reductase inhibitors in follicular growth attenuation. *Phytotherapy Research*, 32(3), 410-418.
- [11] Bodke, V., & Patil, R. (2026). Folklore Herbal Plants as a Safe and Natural Hair Depilatories: A Comprehensive Review. *Journal of Pharmaceutical Research International*, 43(8), 102-111.
- [12] Deshmukh, P., & Shinde, M. (2025). Formulation, Development & Evaluation of Herbal Cream for Depilatory Action. *International Journal of Pharmaceutical Sciences*, 16(2), 224-231.
- [13] Joshi, A., & Chaudhari, V. (2025). Herbal hair removal solution: Exploring plant-based ingredients for safe and natural depilation powder. *International Journal of Cosmetic Science*, 47(4), 315-323.
- [14] More, S., & Tambe, V. (2026). Green cosmeceuticals and global market trends: Shift toward safety-driven botanical cosmetics. *International Cosmeceutical Journal*, 21(1), 45-56.
- [15] Kulkarni, N., & Gawat, S. (2025). Polyherbal cosmetics: Multi-component preservative networks and shelf-life stabilization of sensitive enzymes. *Journal of Pharmaceutical Sciences*, 115(3), 512-520.

