

Applications of Double Aboodh Transform to Boundary Value Problem

I. A. Almardy^{2*}, R. A. Farah^{1,2}, H. Saadoui², K. S. Aboodh¹, A. K. Osman²

Department of Mathematics, Faculty of Science & Technology, Omdurman Islamic University, Khartoum, Sudan¹

Department of Management Information Systems and Production Management

College of Business and Economics, Qassim University, Buraidah, Saudi Arabia²

*Corresponding Author's Email address: imana2627@gmail.com

Abstract: In this paper, we apply the method of the double Aboodh Transform for solving one-dimensional boundary value problems. Through this method, the boundary value problem is solved without converting it into an ordinary differential equation; therefore, there is no need to find the complete solution of an ordinary differential equation. This is the biggest advantage of this method. The main focus of this paper is to develop the method of the double Aboodh transform to solve initial and boundary value problems in applied mathematics.

Keywords: Boundary Value Problem, Double Aboodh Transform, Inverse Aboodh Transform

I. INTRODUCTION

In mathematics, in the field of differential equations, a boundary value problem is a differential equation together with a set of additional constraints, called the boundary conditions [1]. A solution to a boundary value problem is a solution to the differential equation that also satisfies the boundary conditions. Boundary value problems arise in several branches of physics, as any physical differential equation will have them. Problems involving the wave equation, such as the determination of normal modes, are often stated as boundary value problems. To be useful in applications, a boundary value problem should be well posed. This means that given the input to the problem, there exists a unique solution that depends continuously on the input. Much theoretical work in the field of partial differential equations is devoted to proving that boundary value problems arising from scientific and engineering applications are in fact well-posed. Integral transforms are extensively used in solving boundary value problems and integral equations. The problem related to a partial differential equation can be solved by using a special integral transform. Thus, many authors solved the boundary value problems by using a single Laplace. In this study, we use the Double Aboodh Transform to solve the Wave and Heat equation, which is a one-dimensional boundary value problem. Henceforth, the different problems of boundary value are solved without converting them into ordinary differential equations, and there is no need to find a complete solution. So this method is very reliable and convenient for solving boundary value problems. The scheme is put to the test by referring to two different examples.

Some examples of Aboodh Transform from [4][1]:

$f(t)$	$A[f(t)] = F(v)$
	$\frac{1}{v^2}$
t	$\frac{1}{v^3}$
t^2	$\frac{2!}{v^4}$
$t^n, n \in \mathbb{N}$	$\frac{n!}{v^{n+2}}$
e^{at}	$\frac{1}{v^2 - av}$

$\sin(at)$	$\frac{a}{v^3 + a^2v}$
$\cos(at)$	$\frac{1}{v^2 + a^2}$

II. MAIN RESULTS

Example 2.1:

Consider the homogeneous wave equation in the form

$$U_n = c^2 U_{xx}$$

$$U(x, 0) = \sin x, U_t(x, 0) = 2$$

$$U(0, t) = 2t, U_x(0, t) = \cos ct$$

By taking the double Aboodh transform

$$v^2 K(u, v) - K(u, 0) - \frac{1}{v} \frac{\partial K(u, 0)}{\partial t} = c^2 \left[u^2 K(u, v) - K(0, v) - \frac{1}{u} \frac{\partial K(0, v)}{\partial x} \right]$$

The single Aboodh transform of initial conditions gives

$$K(u, 0) = \frac{1}{u(u^2 + 1)}, \frac{\partial K(u, 0)}{\partial t} = \frac{2}{u^2}$$

$$K(0, v) = \frac{2}{v^3}, \frac{\partial K(0, v)}{\partial x} = \frac{1}{v^2 + c^2}$$

Then

$$v^2 K(u, v) - c^2 u^2 K(u, v) = \frac{1}{u(u^2 + 1)} + \frac{2}{u^2 v} - \frac{2c^2}{v^3} - \frac{c^2}{u(v^2 + c^2)}$$

$$(v^2 - c^2 u^2) K(u, v) = \frac{1}{u(u^2 + 1)} + \frac{2}{v} \left(\frac{1}{u^2} - \frac{c^2}{v^2} \right) - \frac{c^2}{u(v^2 + c^2)}$$

$$(v^2 - c^2 u^2) K(u, v) = \frac{1}{u(u^2 + 1)} - \frac{c^2}{u(v^2 + c^2)} + \frac{2}{v} \left(\frac{v^2 - c^2 u^2}{v^2 u^2} \right)$$

$$(v^2 - c^2 u^2) K(u, v) = \frac{v^2 - c^2 u^2}{u(u^2 + 1)(v^2 + c^2)} + \frac{2}{u^2 v^3} (v^2 - c^2 u^2)$$

Then

$$K(u, v) = \frac{1}{u(u^2 + 1)} \frac{1}{(v^2 + c^2)} + \frac{2}{u^2 v^3}$$

Applying inverse double Aboodh transform [1]

$$U(x, t) = \sin x \cos(ct) + 2t$$

Example 2.2:

Solving the heat equation

$$\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2}, t > 0$$

$$U(x, 0) = \sin x,$$

$$U(0, t) = 0, U_x(0, t) = e^{-t}$$

By taking the double Aboodh transform [1] to we get

$$vK(u, v) - u^2 K(u, v) = \frac{1}{v} K(u, 0) - K(0, v) - \frac{1}{u} \frac{\partial K(0, v)}{\partial x}$$

The single Aboodh transform of initial conditions gives

$$K(u, 0) = \frac{1}{u(u^2 + 1)}, \frac{\partial K(0, v)}{\partial x} = \frac{1}{v(v + 1)}, K(0, v) = 0$$

$$(v - u^2)K(u, v) = \frac{1}{v} \left[\frac{1}{u(u^2 + 1)} \right] - \frac{1}{u} \left[\frac{1}{v(v + 1)} \right]$$

$$(v - u^2)K(u, v) = \frac{(v - u^2)}{uv(u^2 + 1)(v + 1)}$$

$$K(u, v) = \frac{1}{uv(u^2 + 1)(v + 1)}$$

Applying inverse double Aboodh transform

$$U(x, t) = e^{-t} \sin x$$

III. CONCLUSION

In this paper, we use double Aboodh Transform and the standard properties of Aboodh Transform are discussed for solving one-dimensional boundary value problems.

IV. ACKNOWLEDGEMENTS

The authors extend their appreciation to the Deputyship for Research & Innovation, Ministry of Education and Qassim University, Saudi Arabia for funding this research work through the project number (QU- 6640- 51452).

REFERENCES

- [1]. K. S. Aboodh, R. A. Farah, I. A. Almardy and F. A. Almostafa, Solution of partial Integro-Differential Equations by using Aboodh and Double Aboodh Transform Methods, Global Journal of pure and Applied Mathematics, ISSN 0973-1768 Volume 13, Number 8 (2017), pp.4347-4360.
- [2]. A. M. Wazwaz, A reliable modification of Adomian's decomposition method, Appl. Math. And Comput., 92(1)(1998), 1-7.
- [3]. D. G. Duff, Transform Methods for solving Partial Differential Equations, Chapman and Hall/CRC, Boca Raton, F. L., (2004).
- [4]. K. S. Aboodh, R. A. Farah, I. A. Almardy and F. A. Almostafa, some Application of Aboodh Transform to First Order Constant Coefficients Complex equations, International Journal of Mathematics and its Applications, ISSN : 2347-1557, App.6(1-A)(2018), 1-6.
- [5]. Adem Kilicman and Hassan Eltayeb, A note on defining singular integral as distribution and partial differential equations with convolution term, Mathematical & Computer Modelling, 49(2013), 327-336.
- [6]. A. Estrin and T. J. Higgins, The Solution of Boundary Value Problems by Multiple Laplace Transformation, Journal of the Franklin Institute, 252(2)(1951), 153-167.
- [7]. K. S. Aboodh, M. Y. Ahmed, R.A. Farah, I. A. Almardy and M. Belkhamza, New Transform Iterative Method for Solving some Klein-Gordon Equations, (IJARSCT) IIUI, ISSN 1 Volume 2, (2022), pp.118-126. SCOPe Database Article Link: <https://sdindex.com/documents/00000310/00001-85016.pdf>
- [8]. K. S. Aboodh, I. A. Almardy, R. A. Farah, M. Y. Ahmed and R. I. Nuruddeen, On the Application of Aboodh Transform to System of Partial Differential Equations, BEST, IJHAMS Journal, ISSN(P): 2348-0521; ISSN(E): 2454-4728 Volume 10, Issue 2, Dec 2022.